

1. Essential drugs are
 - a. Life saving drugs
 - b. Drugs that meet the priority health care needs of the population
 - c. Drugs that are used in medical emergencies
 - d. Drugs that are listed in the pharmacopoeia of a country
2. The loading dose of the drug is governed by its
 - a. Renal clearance
 - b. Plasma half life
 - c. Volume of distribution
 - d. Elimination rate constant
3. Pseudocholinesterase differ from true cholinesterase in that
 - a. It hydrolyses acetylcholine at slower rate
 - b. It is more susceptible to inhibition by physostigmine
 - c. It doesn't hydrolyse acetylcholine
 - d. It is the only form of circulating cholinesterase
4. Bromhexine acts by
 - a. Inhibiting cough centre
 - b. Depolymerizing mucopolysaccharides present in sputum
 - c. Irritating gastric mucosa and reflexly increasing bronchial secretion
 - d. Desensitizing stretch receptors in lung
5. The following general anesthetic has poor muscle relaxation
 - a. Ether
 - b. Halothane
 - c. Nitrous oxide
 - d. Sevoflurane
6. The antiepileptic drug that is effective in manic depressive illness also is
 - a. Phenobarbitone
 - b. Ethosuximide
 - c. Phenytoin
 - d. Carbamazepine
7. The major limitation to the use of clozapine for treatment of schizophrenia is
 - a. Its inability to benefit negative symptoms of schizophrenia
 - b. Its potential to cause agranulocytosis
 - c. It produces hyperprolactinemia
 - d. High incidence of extra pyramidal symptoms
8. Select the most suitable antiarrhythmic drug for ventricular arrhythmia due to digoxin toxicity
 - a. Lidocaine
 - b. Propranolol
 - c. Verapamil
 - d. Quinidine

9. Glyceryl nitrate is administered by following routes except
a. Intravenous b. Intramuscular c. Oral d. Sublingual
10. Reversible loss of taste sensation can be a side effect of the following anti hypertensive drug
a. Prazosin b. Verapamil c. Propranolol d. Captopril
11. The following drug is a poor inhibitor of prostaglandin synthesis in peripheral tissues but more active in brain
a. Mefenamic acid b. Ketorolac c. Paracetamol d. Nimesulide
12. Prolonged lithium therapy can cause
a. Parkinsonism b. Goiter c. Diabetes mellitus d. Gout
13. Which of the following drug is a potassium channel opener?
a. Nicorandil b. Hydralazine c. Amiloride d. Glibenclamide
14. Indication of desmopressin include the following except
a. Nephrogenic diabetes insipidus b. Neurogenic diabetes insipidus
c. Bleeding due to hemophilia d. Bedwetting in children
15. The following is not a valid indication for parenteral iron therapy
a. Inadequate response to oral iron due to patient non compliance
b. Anemia during pregnancy
c. Anemia in a patient with active rheumatoid arthritis
d. Severe anemia associated with chronic bleeding
16. The primary mechanism by which heparin inhibits coagulation of blood is
a. Inhibition of factor XII a and XIII a
b. Activation of antithrombin III with inhibition of factors IX and XI
c. Inhibition of factor Xa and thrombin mediated conversion of fibrinogen to fibrin
d. Direct inhibition of prothrombin to thrombin conversion
17. Select the fibrinolytic drug that is antigenic
a. Streptokinase b. Urokinase c. Alteplase d. All
18. Which of the following hypolipidemic drug is most effective in increasing HDL cholesterol level and lowering serum triglyceride?
a. Cholestyramine b. Fenofibrate c. Pravastatin d. Nicotinic acid
19. Tranexemic acid is used in following condition
a. Organophosphorus compound poisoning b. Barbiturate poisoning
c. Excessive bleeding due to fibrinolytic drugs d. Bleeding due to heparin overdose
20. The following drug is neither analgesic nor anti-inflammatory and do not have uricosuric effect, but highly efficacious in acute gout
a. Naproxen b. Colchicine c. Prednisolone d. Sulfapyrazone

THE TAMIL NADU DR. M.G.R. MEDICAL UNIVERSITY

[MBBS 0822]

AUGUST 2022

Sub. Code :6061

M.B.B.S. DEGREE EXAMINATION

(For the candidates admitted from the Academic Year 2019-2020)

SECOND YEAR – SUPPLEMENTARY (CBME)

PAPER I – PHARMACOLOGY – I

Q.P. Code: 526061

Time: 30 Minutes

Maximum : 20 Marks

Answer All Questions

**Choose one correct answer in the box provided in the Answer Script.
No overwriting should be done.**

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Which of the following is not a Cytochrome P450 enzyme inducer
a. Phenytoin b. Phenobarbitone c. Erythromycin d. Cigarette smoke
2. Drugs with high Plasma Protein Binding have
a. Short duration of action b. Less drug interactions
c. Lower Volume of distribution d. all the above
3. Which is a prodrug?
a. Ampicillin b. Captopril c. Levodopa d. Phenytoin
4. Anti-hypertensive drug causing erectile dysfunction
a. Calcium channel blocker b. ACE inhibitor
c. AT1 receptor antagonists d. β -blockers
5. Pilocarpine induces
a. Active miosis b. Passive miosis c. Active mydriasis d. Passive mydriasis
6. Propranolol can be used in all except
a. Thyrotoxicosis b. Variant angina c. Migraine d. HT
7. Fenoldopam is used in the management of
a. Hypertensive emergencies b. Congestive heart failure
c. Migraine prophylaxis d. Tachyarrhythmia
8. Mechanism of action of digitalis is
a. Inhibits Na^+K^+ ATPase pump b. Inhibits Na^+H^+ ATPase pump
c. Activates Na^+Cl^- pump d. Inhibits carbonic anhydrase
9. Bilateral renal artery stenosis is a contraindication for which Antihypertensive drug?
a. Beta-blockers b. Calcium channel blockers
c. ACE inhibitors d. Diuretics

10. All of the following are CNS stimulants except
a. Amphetamines b. Benzodiazepines
c. Cocaine d. Methylphenidate
11. Which of the following drug is used as Transcranial patch for Parkinsons disease?
a. Rotigotine b. Levodopa c. Selegiline d. carbidopa
12. The side effect of Phenytoin when its plasma concentration is above therapeutic level is
a. Gum hypertrophy b. Hirsutism C. Osteomalacia d. Ataxia
13. Which statement is true about Carbamazepine?
a. Used in Trigeminal neuralgia
b. Carbamazepine is an enzyme inhibitor
c. Can cause megaloblastic anaemia
d. Drug of choice for Status epilepticus
14. Which one of the following drugs is contraindicated in acute myocardial infarction.
a. Morphine b. Pentazocine c. Nitroglycerin d. Beta Blockers
15. Anti-craving agents for alcohol dependence are all except
a. Lorazepam b. Acamprosate c. Topiramate d. Naltrexone
16. What is the drug of choice for obsessive compulsive disorder?
a. Imipramine b. Fluoxetine c. Benzodiazepines d. Alprazolam
17. Leukotriene receptor antagonist used for bronchial asthma is
a. Zileuton b. Cromolyn sodium c. Zafirlukast d. Aminophylline
18. Preconceptional intake of which reduces the incidence of neural tube defects?
a. Folic acid b. Vitamin A c. Vitamin C d. Vitamin E
19. Filgrastim is used for treatment of
a. Neutropenia b. Anaemia c. Polycythaemia d. Neutrophilia
20. Absolute contraindication to thrombolytic therapy is
a. Pregnancy b. History of haemorrhagic stroke in past one year
c. Patient on nitrates d. hypertension.

[MBBS 0822]

THE TAMIL NADU DR. M.G.R. MEDICAL UNIVERSITY

[MBBS 0223]

FEBRUARY 2023

Sub. Code :6061

M.B.B.S. DEGREE EXAMINATION

(For the candidates admitted from the Academic Year 2019-2020)

SECOND YEAR – (CBME)

PAPER I – PHARMACOLOGY – I

Q.P. Code: 526061

Time: 30 Minutes

Maximum : 20 Marks

Answer All Questions

Choose one correct answer in the box provided in the Answer Script. No overwriting should be done. Choice should be given in Capital Letters.

III. Multiple Choice Questions:

(20 x 1 = 20)

1. A patient with iron deficiency anaemia is put on iron therapy. What should be the rate of rise of Hb level so that response is considered adequate?
A) 0.05 to 0.1g/dl per week B) 0.1 to 0.2 g/ dl per week
C) more than 1g/ dl per week D) 0.5 to 1g / dl per week
2. A patient with coronary artery disease has raised triglyceride levels but normal cholesterol level. Which hypolipidemic drug should be prescribed?
A) Lovastatin B) Gemfibrozil C) Cholestyramine D) Nicotinic acid
3. Marked redistribution is a feature of
A) Highly lipid soluble drugs B) Poorly lipid soluble drugs
C) Depot preparation D) High plasma protein bound drugs
4. The CYP450 isoenzyme involved in metabolism of largest number of drugs in human beings has been implicated in some dangerous drug interactions
A) CYP1A2 B) CYP2E1 C) CYP2C9 D) CYP3A4
5. The receptor that is located intracellularly
A) Opioid mu receptor B) Steroid receptor
C) Angiotensin receptor D) Prostaglandin receptor
6. The calcium channel blocker used for prophylaxis of migraine but not angina pectoris
A) Verapamil B) Amlodipine C) Flunarizine D) Diltiazem
7. Which of the following is nonsedative anxiolytic?
A) Alprazolam B) Hydroxyzine C) Chlorpromazine D) Buspirone
8. The drug that release histamine from mast cell without involving in immunological reaction is
A) pirin B) Morphine C) Procaine D) Erythromycin
9. The following is a selective 5HT 1B/1D Receptor agonist
A) Ondansetron B) Sumatriptan C) Cyproheptadine D) Risperidone

10. Low dose aspirin prolong bleeding time by selectively inhibiting following mediator in platelets
A) Thromboxane A2 B) Prostacyclin C) Histamine D) 5HT
11. The dose limiting adverse effect of colchicine is
A) Renal damage B) Muscular paralysis C) Sedation D) Diarrhoea
12. Higher dose of theophylline is used to maintain therapeutic plasma concentration in
A) Those receiving cimetidine B) Those receiving erythromycin
C) Smokers D) Chronic renal failure patients
13. Postoperative muscle soreness and myalgia may be side effects of following neuromuscular blockers
A) Vecuronium B) Atracurium C) Succinylcholine D) Tubocurarine
14. Which one of the following is an example for Physiological Antagonism
A) Acetylcholine and Atropine B) Adrenaline and Histamine
C) Heparin and Protamine sulphate D) Acetylcholine and physostigmine
15. Ventricular remodeling after myocardial infarction involves mediation of
A) Angiotensin II B) Prostaglandin C) Thromboxane A2 D) Bradykinin
16. The diuretic preferred for mobilising edema fluid in congestive cardiac failure is
A) Hydrochlorothiazide B) Amiloride C) Metolazone D) Furosemide
17. The following antiarrhythmic drug has prominent anticholinergic action
A) Lignocaine B) Disopyramide C) Amiodarone D) Quinidine
18. Select the drug which is potassium channel opener as well as nitric oxide donor
A) Diazoxide B) Minoxidil C) Nicorandil D) Sodium nitroprusside
19. Inhaled salbutamol is useful in bronchial asthma for
A) Round the clock prophylaxis of asthma B) Status asthmatics
C) Aborting / terminating asthma attacks D) All of the above
20. The following antihypertensive drug is topically used to treat alopecia areata
A) Minoxidil B) Hydralazine C) Clonidine D) Nifedipine.

[MBBS 0223]

THE TAMIL NADU DR. M.G.R. MEDICAL UNIVERSITY

[MBBS 0723]

JULY 2023

Sub. Code :6061

M.B.B.S. DEGREE EXAMINATION

(For the candidates admitted from the Academic Year 2019-2020)

SECOND YEAR – (CBME)

PAPER I – PHARMACOLOGY – I

Q.P. Code: 526061

Time: 30 Minutes

Maximum : 20 Marks

Answer All Questions

Choose one correct answer in the box provided in the Answer Script. No overwriting should be done. Choice should be given in Capital Letters.

III. Multiple Choice Questions: (20 x 1 = 20)

1. Choose a drug that has long plasma half life
A) Dopamine B) Alteplase C) Chloroquine D) Esmolol
2. Therapeutic drug monitoring is required for
A) Drugs with Low safety margin B) Hit and run drugs
C) Drugs whose response is measurable D) Drugs that develop tolerance
3. Selective Beta₂ stimulant used as uterine relaxant is
A) Ephedrine B) Brimonidine C) Clonidine D) Isoxsuprine
4. Most preferred drug in Chronic congestive cardiac failure is
A) Dopamine B) Enalapril C) Prazosin D) Nifedepine
5. First line drug in open angle glaucoma is
A) Latanoprost B) Pilocarpine C) Acetazolamide D) Dipivefrine
6. One of the following is a contraindication to Beta blockers
A) Cirrhosis B) Alcohol withdrawal
C) Heart block D) Esophageal varices
7. Atropine substitute preferred in Bronchial asthma is
A) Glycopyrrolate B) Ipratropium C) Tropicamide D) Benztropine
8. Coronary steal phenomenon is seen with
A) Dipyridamole B) Diltiazem C) Propranolol D) Verapamil
9. Sumatriptan is an agonist at which serotonergic receptor
A) 5-HT_{1A} B) 5-HT_{1D} C) 5-HT_{2A} D) 5HT_{2C}
10. Uricosuric drug used to treat chronic gout among the following is
A) Allopurinol B) Cochicine C) Febuxostat D) Probenecid

11. More cardiotoxic local anaesthetics is
A) Bupivacaine B) Lignocaine C) Prilocaine D) Procaine
12. Flumazenil is the antagonist of
A) Phenobarbitone B) Diazepam C) Morphine D) Phenytoin
13. The orally active H₁ Histamine analogue used in vertigo is
A) Betahistine B) Diphenhydramine
C) Clemastine D) Rupatadine
14. The preferred general anaesthetic in children is
A) Thiopentone sodium B) Isoflurane C) Sevoflurane D) Propofol
15. Typical anti-psychotic less prone for extra pyramidal side effect
A) Chlorpromazine B) Thioridazine C) Fluphenazine D) Trifluoperazine
16. Digoxin acts by inhibiting
A) H⁺ K⁺ ATPase B) Na⁺ K⁺ 2Cl⁻ co-transport
C) Na⁺ Cl⁻ symport D) Na⁺ K⁺ ATPase
17. Dabigatran is
A) Direct factor Xa inhibitor B) Thromboxane synthesis inhibitor
C) Direct thrombin inhibitor D) Phosphodiesterase inhibitor
18. Contraindication for use of ACE inhibitor
A) Bilateral renal artery stenosis B) Diabetic nephropathy
C) Left ventricular hypertrophy D) Congestive cardiac failure
19. Drug of choice in lithium induced nephrogenic diabetes insipidus
A) Furosemide B) Acetazolamide C) Amiloride D) Mannitol
20. Drug of choice in malignant hyperthermia
A) Baclofen B) Dantrolene C) Tizanidine D) Chlorzoxazone.

[MBBS 0723]

THE TAMIL NADU DR. M.G.R. MEDICAL UNIVERSITY

[MBBS 0224]

FEBRUARY 2024

Sub. Code :6061

M.B.B.S. DEGREE EXAMINATION

(For the candidates admitted from the Academic Year 2019-2020)

SECOND PROFESSIONAL YEAR – (CBME)

PAPER I – PHARMACOLOGY

Q.P. Code: 526061

Time: 30 Minutes

Maximum : 20 Marks

Answer All Questions

**Choose one correct answer in the box provided in the Answer Script.
No overwriting should be done. Choice should be given in Capital Letters.**

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Essential drugs are
 - A) Life saving drugs
 - B) Drugs that meet the priority health care needs of the population
 - C) Drugs that must be present in the emergency bag of a doctor
 - D) Drugs that are listed in the pharmacopoeias of a country
2. The elderly patients are relatively intolerant to:
 - A) Digoxin
 - B) Salbutamol
 - C) Propranolol
 - D) Nifedipine
3. 3 Years old child undergoing eye surgery had initial heart rate of 140 beats/min. After anaesthesia and short surgery, heart rate dropped to 40 beats/min. Whats should be the next step?
 - A) Stop surgery
 - B) Decrease plane of anaesthesia
 - C) Inj. Glycopyrrolate
 - D) Inj. Atropine
4. Preferred Skeletal muscle relaxants for patients with Liver/Kidney disease & elderly?
 - A) d-TC
 - B) Pancuronium
 - C) Cisatracurium
 - D) Pipecuronium
5. Which of the following routes allows titration of dose of a drug with its response of administration?
 - A) Intravenous
 - B) Transdermal
 - C) Intramuscular
 - D) Subcutaneous
6. The drug of choice for hyperkinetic children is:
 - A) Methylphenidate
 - B) Nikethamide
 - C) Caffeine
 - D) Clonazepam
7. Preferred drug for medical closure of Patent Ductus Arteriosus?
 - A) Aspirin
 - B) Nefopam
 - C) Ibuprofen
 - D) Indomethacin
8. Which direct thrombin inhibitor produces hepatotoxicity?
 - A) Hirudin
 - B) Lepirudin
 - C) Argatroban
 - D) Ximelagatran
9. Recombinant human erythropoietin is indicated for
 - A) Megaloblastic anaemia
 - B) Haemolytic anaemia
 - C) Anaemia in patients of thalassemia
 - D) Anaemia in chronic renal failure patients

10. Sargramostim is a growth factor of?
A) Megakaryocyte growth factor B) GM - CSF
C) G - CSF D) Erythropoietin
11. Which of the drug is used by intravenous infusion for emergency control of tachycardia and sudden rise in blood pressure?
A) Amiodarone B) Lignocaine C) Esmolol D) Disopyramide
12. Which of the following peptides is a selective vasopressin V2 receptor agonist?
A) Arginine vasopressin B) Desmopressin
C) Lypressin D) Terlipressin
13. Absorption of oral iron preparations can be facilitated by co-administering
A) Antacids B) Tetracyclines C) Phosphates D) Ascorbic acid
14. In cancer/ palliative care patients on opioid analgesics- c/o constipation; suggest a drug used to relieve constipation without reducing analgesic effect
A) Pentazocine B) Nalbuphine
C) Methyl naltrexone D) Buprenorphine
15. 1st choice of drug used in treatment of chronic gout?
A) Colchicine B) Allopurinol C) Probenecid D) Pegloticase
16. 1st line of drug for Migraine patient those who doesn't respond to analgesics
A) TCA B) Ca⁺ channel blockers
C) Selective 5HT Agonists/ Ergot D) β - blockers
17. Loperamide acts by?
A) Δ opioid agonist action B) κ opioid agonist action
C) μ opioid agonist action D) None
18. In cardiac cells, procainamide
A) Decreased automaticity B) Increases refractory period
C) Slows down conduction D) All of the above
19. The rabbit syndrome is treated by :
A) Lorazepam B) Lithium
C) Levodopa D) Trihexyphenidyl
20. Identify correct anti-inflammatory dose of Aspirin
A) 75-162 mg/d B) 3-6 gm/d
C) 0.3-0.6 gm/d D) 30-60 mg/d

[MBBS 0224]

THE TAMIL NADU DR. M.G.R. MEDICAL UNIVERSITY

[MBBS 0524]

MAY 2024

Sub. Code :6061

M.B.B.S. DEGREE EXAMINATIONS

(For the candidates admitted from the Academic Year 2019-2020)

SECOND PROFESSIONAL – SUPPLEMENTARY (CBME)

PAPER I – PHARMACOLOGY

Q.P. Code: 526061

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

**Choose one correct answer in the box provided in the Answer Script.
No overwriting should be done. Choice should be given in Capital Letters.**

III. Multiple Choice Questions: (20 x 1 = 20)

1. Which one is not a G-Protein coupled Receptor transduction pathway
A) Ligand gated ion channels B) Adenyl cyclase pathway
C) Phospholipase pathway D) Channel regulation
2. Genetically determined disease is
A) Side effect B) Toxic effect C) Idiosyncrasy D) Secondary effect
3. ALPHA-2 Selective antagonist is
A) Terazosin B) Chlorpromazine C) Dihydroergotamine D) Yohimbine
4. The therapeutic agent that increase the uveoscleral outflow
A) Pilocarpine B) Acetazolamide C) Latanoprost D) Timolol
5. The muscarinic (M-3) receptor antagonist is
A) Pirenzepine B) Solifenacin C) Methacholine D) Glycopyrrolate
6. The L -Type of Calcium Channel blocker is
A) Flunarizine B) Ethosuxamide C) Mibefradil D) Verapamil
7. The following drug is not a Phosphodiesterase-3 inhibitor is
A) Amrinone B) Ivabradine C) Levosimendan D) Milrinone
8. Which one of the following is not a adverse effect of Endothelin Receptor Antagonist
A) Anemia B) Increased amino transferase level
C) Teratogenicity D) Neuropathy
9. Adrenergic Neuron blocker is
A) Guanethidine B) Hydralazine C) Methyl dopa D) Clonidine
10. The drug which is not highly bound to ALPHA-1acid glycoprotein is
A) Lidocaine B) Quinidine C) Verapamil D) Warfarin
11. Choose the drug used to maintain the patency of ductus arteriosus
A) Dinoprostone B) Carboprost C) Alprostadil D) Misoprostol

12. Paracetamol toxicity occurs due to the following Metabolite:
A) Hepatic glutathione B) N-Acetyl P-Benzoquinone imine
C) Nortriptyline D) N acetyl Procainamide
13. The local anaesthetic agent used for oesophagus, stomach site is
A) Tetracaine B) Lidocaine C) Cocaine D) Oxethazine
14. Selective serotonin receptor antagonist is
A) Doxepin B) Nortriptyline C) Sertraline D) Reboxetine
15. Opioid Receptor antagonist used in the treatment of postoperative ileus is
A) Nalmefene B) Nalbuphine C) Naloxone D) Methylnaltrexone
16. The cognition enhancer which has cholinergic activator property is
A) Citicoline B) Donepezil C) Piribedil D) Memantine
17. Dissociative anesthesia is caused by
A) Ketamine B) Diazepam C) Thio pentone D) Fentanyl
18. Vasopressin antagonists is
A) Chlopropamide B) Tolvaptan C) Indomethacin D) Terlipressin
19. Nonopioid Antitussives is
A) Noscapine B) Ethylmorphine C) Ambroxol D) Pholcodeine
20. In zero order kinetics
A) Clearance remain constant
B) Constant fraction of the drug is eliminated
C) Clearance decreases with increase in concentration
D) Apply to the drugs which do not saturate the elimination

[MBBS 0524]

M.B.B.S. DEGREE EXAMINATION

(For the candidates admitted from the Academic Year 2019-2020)

SECOND PROFESSIONAL – MBBS (CBME)

PAPER I – PHARMACOLOGY

Q.P. Code: 526061

SET - A

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Alkalinization of urine hastens the excretion of:
A) Weakly basic drugs B) Weakly acidic drugs
C) Strongly acidic drugs D) Nonpolar drugs
2. If the total amount of a drug present in the body at a given moment is 2.0 g and its plasma concentration is 50 µg/ml, its volume of distribution is:
A) 100 L B) 40 L C) 60 L D) 50 L
3. Induction of drug metabolizing enzymes involves:
A) A conformational change in the enzyme protein to favour binding of substrate molecules
B) Expression of enzyme molecules on the surface of hepatocytes
C) Enhanced transport of substrate molecules into hepatocytes
D) Increased synthesis of enzyme protein
4. Receptor agonists possess:
A) Affinity but no intrinsic activity
B) Intrinsic activity but no affinity
C) Affinity and intrinsic activity with a + sign
D) Affinity and intrinsic activity with a – sign
5. The choline ester resistant to both true and pseudocholinesterase is:
A) Methacholine B) Bethanechol C) Benzoylcholine D) Butyrylcholine
6. Sumatriptan acts by the following mechanism
A) 5HT 1D antagonist B) 5HT 1D/B agonist
C) 5HT 3 antagonist D) 5HT 1D/B antagonist
7. The principal process which terminates the action of noradrenaline released from adrenergic nerve ending is:
A) Degradation by MAO B) Methylation by COMT
C) Axonal uptake D) Extra neuronal uptake
8. Select the H1 antihistaminic which is used topically in the nose for allergic rhinitis:
A) Loratadine B) Cetirizine C) Fexofenadine D) Azelastine
9. Aspirin is contraindicated in pregnant women near term because:
A) Labour may be delayed and prolonged
B) Blood loss during delivery may be more
C) Foetus may suffer premature closure of ductus arteriosus
D) All of the above

10. Bronchodilators are useful in cough:
A) Only when cough is nonproductive
B) Only when cough is associated with thick sticky secretions
C) Only when reflex bronchoconstriction is associated
D) Irrespective of nature of cough or associated features
11. No surgical operation should be performed during the following stage of anaesthesia:
A) Stage I B) Stage II C) Stage III, plane 1 D) Stage III, plane 3
12. Benzodiazepines differ from barbiturates in the following aspects except:
A) They have a steeper dose response curve
B) They have higher therapeutic index
C) They have lower abuse liability
D) They do not induce microsomal drug metabolizing enzymes
13. The following drug combination should not be used in parkinsonism:
A) Levodopa + anticholinergics B) Levodopa + amantadine
C) Bromocriptine + levodopa D) Amantadine + anticholinergics
14. The following side effect of neuroleptics responds to propranolol:
A) Parkinsonism B) Acute muscle dystonia
C) Tardive dyskinesia D) Akathisia
15. Limitations of typical tricyclic anti-depressants include the following except:
A) Narrow safety margin B) Low oral bioavailability
C) Frequent side effects D) Long latent period for response
16. Digoxin is contraindicated in:
A) Angina pectoris patients B) Ventricular tachycardia
C) Hypertensive patients D) Partial heart-block
17. The following is an orally active lignocaine congener used for both acute as well as chronic ventricular arrhythmias:
A) Mexiletine B) Flecainide C) Prilocaine D) Propafenone
18. The following calcium channel blocker is specifically indicated to counteract cerebral vasospasm and neurological sequelae following subarachnoid haemorrhage:
A) Lacidipine B) Nimodipine C) Nicardipine D) Nitrendipine
19. Use of potassium sparing diuretics in patients receiving the following drug needs close monitoring:
A) Furosemide B) Hydrochlorothiazide C) Captopril D) Verapamil
20. The following statements are true of oral anticoagulants except:
A) They interfere with an early step in the synthesis of clotting factors
B) Irrespective of the dose administered, their anticoagulant effect has a latency of onset of 1-3 days
C) Their dose is adjusted by repeated measurement of prothrombin time
D) They are contraindicated during pregnancy

M.B.B.S. DEGREE EXAMINATION

(For the candidates admitted from the Academic Year 2019-2020)

SECOND PROFESSIONAL – MBBS (CBME)

PAPER I – PHARMACOLOGY

Q.P. Code: 526061

SET - B

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Sumatriptan acts by the following mechanism
A) 5HT 1D antagonist B) 5HT 1D/B agonist
C) 5HT 3 antagonist D) 5HT 1D/B antagonist
2. The principal process which terminates the action of noradrenaline released from adrenergic nerve ending is:
A) Degradation by MAO B) Methylation by COMT
C) Axonal uptake D) Extra neuronal uptake
3. Select the H1 antihistaminic which is used topically in the nose for allergic rhinitis:
A) Loratadine B) Cetirizine C) Fexofenadine D) Azelastine
4. Aspirin is contraindicated in pregnant women near term because:
A) Labour may be delayed and prolonged
B) Blood loss during delivery may be more
C) Foetus may suffer premature closure of ductus arteriosus
D) All of the above
5. Bronchodilators are useful in cough:
A) Only when cough is nonproductive
B) Only when cough is associated with thick sticky secretions
C) Only when reflex bronchoconstriction is associated
D) Irrespective of nature of cough or associated features
6. No surgical operation should be performed during the following stage of anaesthesia:
A) Stage I B) Stage II C) Stage III, plane 1 D) Stage III, plane 3
7. Benzodiazepines differ from barbiturates in the following aspects except:
A) They have a steeper dose response curve
B) They have higher therapeutic index
C) They have lower abuse liability
D) They do not induce microsomal drug metabolizing enzymes
8. The following drug combination should not be used in parkinsonism:
A) Levodopa + anticholinergics B) Levodopa + amantadine
C) Bromocriptine + levodopa D) Amantadine + anticholinergics
9. The following side effect of neuroleptics responds to propranolol:
A) Parkinsonism B) Acute muscle dystonia
C) Tardive dyskinesia D) Akathisia

10. Limitations of typical tricyclic anti-depressants include the following except:
A) Narrow safety margin B) Low oral bioavailability
C) Frequent side effects D) Long latent period for response
11. Digoxin is contraindicated in:
A) Angina pectoris patients B) Ventricular tachycardia
C) Hypertensive patients D) Partial heart-block
12. The following is an orally active lignocaine congener used for both acute as well as chronic ventricular arrhythmias:
A) Mexiletine B) Flecainide C) Prilocaine D) Propafenone
13. The following calcium channel blocker is specifically indicated to counteract cerebral vasospasm and neurological sequelae following subarachnoid haemorrhage:
A) Lacidipine B) Nimodipine C) Nicardipine D) Nitrendipine
14. Use of potassium sparing diuretics in patients receiving the following drug needs close monitoring:
A) Furosemide B) Hydrochlorothiazide C) Captopril D) Verapamil
15. The following statements are true of oral anticoagulants except:
A) They interfere with an early step in the synthesis of clotting factors
B) Irrespective of the dose administered, their anticoagulant effect has a latency of onset of 1-3 days
C) Their dose is adjusted by repeated measurement of prothrombin time
D) They are contraindicated during pregnancy
16. Alkalinization of urine hastens the excretion of:
A) Weakly basic drugs B) Weakly acidic drugs
C) Strongly acidic drugs D) Nonpolar drugs
17. If the total amount of a drug present in the body at a given moment is 2.0 g and its plasma concentration is 50 µg/ml, its volume of distribution is:
A) 100 L B) 40 L C) 60 L D) 50 L
18. Induction of drug metabolizing enzymes involves:
A) A conformational change in the enzyme protein to favour binding of substrate molecules
B) Expression of enzyme molecules on the surface of hepatocytes
C) Enhanced transport of substrate molecules into hepatocytes
D) Increased synthesis of enzyme protein
19. Receptor agonists possess:
A) Affinity but no intrinsic activity
B) Intrinsic activity but no affinity
C) Affinity and intrinsic activity with a + sign
D) Affinity and intrinsic activity with a – sign
20. The choline ester resistant to both true and pseudocholinesterase is:
A) Methacholine B) Bethanechol C) Benzoylcholine D) Butyrylcholine

M.B.B.S. DEGREE EXAMINATION

(For the candidates admitted from the Academic Year 2019-2020)

SECOND PROFESSIONAL – MBBS (CBME)

PAPER I – PHARMACOLOGY

Q.P. Code: 526061

SET - C

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

III. Multiple Choice Questions:

(20 x 1 = 20)

1. No surgical operation should be performed during the following stage of anaesthesia:
A) Stage I B) Stage II C) Stage III, plane 1 D) Stage III, plane 3
2. Benzodiazepines differ from barbiturates in the following aspects except:
A) They have a steeper dose response curve
B) They have higher therapeutic index
C) They have lower abuse liability
D) They do not induce microsomal drug metabolizing enzymes
3. The following drug combination should not be used in parkinsonism:
A) Levodopa + anticholinergics B) Levodopa + amantadine
C) Bromocriptine + levodopa D) Amantadine + anticholinergics
4. The following side effect of neuroleptics responds to propranolol:
A) Parkinsonism B) Acute muscle dystonia
C) Tardive dyskinesia D) Akathisia
5. Limitations of typical tricyclic anti-depressants include the following except:
A) Narrow safety margin B) Low oral bioavailability
C) Frequent side effects D) Long latent period for response
6. Digoxin is contraindicated in:
A) Angina pectoris patients B) Ventricular tachycardia
C) Hypertensive patients D) Partial heart-block
7. The following is an orally active lignocaine congener used for both acute as well as chronic ventricular arrhythmias:
A) Mexiletine B) Flecainide C) Prilocaine D) Propafenone
8. The following calcium channel blocker is specifically indicated to counteract cerebral vasospasm and neurological sequelae following subarachnoid haemorrhage:
A) Lacidipine B) Nimodipine C) Nicardipine D) Nitrendipine
9. Use of potassium sparing diuretics in patients receiving the following drug needs close monitoring:
A) Furosemide B) Hydrochlorothiazide C) Captopril D) Verapamil

10. The following statements are true of oral anticoagulants except:
 - A) They interfere with an early step in the synthesis of clotting factors
 - B) Irrespective of the dose administered, their anticoagulant effect has a latency of onset of 1-3 days
 - C) Their dose is adjusted by repeated measurement of prothrombin time
 - D) They are contraindicated during pregnancy
11. Alkalinization of urine hastens the excretion of:
 - A) Weakly basic drugs
 - B) Weakly acidic drugs
 - C) Strongly acidic drugs
 - D) Nonpolar drugs
12. If the total amount of a drug present in the body at a given moment is 2.0 g and its plasma concentration is 50 µg/ml, its volume of distribution is:
 - A) 100 L
 - B) 40 L
 - C) 60 L
 - D) 50 L
13. Induction of drug metabolizing enzymes involves:
 - A) A conformational change in the enzyme protein to favour binding of substrate molecules
 - B) Expression of enzyme molecules on the surface of hepatocytes
 - C) Enhanced transport of substrate molecules into hepatocytes
 - D) Increased synthesis of enzyme protein
14. Receptor agonists possess:
 - A) Affinity but no intrinsic activity
 - B) Intrinsic activity but no affinity
 - C) Affinity and intrinsic activity with a + sign
 - D) Affinity and intrinsic activity with a – sign
15. The choline ester resistant to both true and pseudocholinesterase is:
 - A) Methacholine
 - B) Bethanechol
 - C) Benzoylcholine
 - D) Butyrylcholine
16. Sumatriptan acts by the following mechanism
 - A) 5HT 1D antagonist
 - B) 5HT 1D/B agonist
 - C) 5HT 3 antagonist
 - D) 5HT 1D/B antagonist
17. The principal process which terminates the action of noradrenaline released from adrenergic nerve ending is:
 - A) Degradation by MAO
 - B) Methylation by COMT
 - C) Axonal uptake
 - D) Extra neuronal uptake
18. Select the H₁ antihistaminic which is used topically in the nose for allergic rhinitis:
 - A) Loratadine
 - B) Cetirizine
 - C) Fexofenadine
 - D) Azelastine
19. Aspirin is contraindicated in pregnant women near term because:
 - A) Labour may be delayed and prolonged
 - B) Blood loss during delivery may be more
 - C) Foetus may suffer premature closure of ductus arteriosus
 - D) All of the above
20. Bronchodilators are useful in cough:
 - A) Only when cough is nonproductive
 - B) Only when cough is associated with thick sticky secretions
 - C) Only when reflex bronchoconstriction is associated
 - D) Irrespective of nature of cough or associated features

M.B.B.S. DEGREE EXAMINATION

(For the candidates admitted from the Academic Year 2019-2020)

SECOND PROFESSIONAL – MBBS (CBME)

PAPER I – PHARMACOLOGY

Q.P. Code: 526061

SET - D

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Digoxin is contraindicated in:
A) Angina pectoris patients B) Ventricular tachycardia
C) Hypertensive patients D) Partial heart-block
2. The following is an orally active lignocaine congener used for both acute as well as chronic ventricular arrhythmias:
A) Mexiletine B) Flecainide C) Prilocaine D) Propafenone
3. The following calcium channel blocker is specifically indicated to counteract cerebral vasospasm and neurological sequelae following subarachnoid haemorrhage:
A) Lacidipine B) Nimodipine C) Nicardipine D) Nitrendipine
4. Use of potassium sparing diuretics in patients receiving the following drug needs close monitoring:
A) Furosemide B) Hydrochlorothiazide C) Captopril D) Verapamil
5. The following statements are true of oral anticoagulants except:
A) They interfere with an early step in the synthesis of clotting factors
B) Irrespective of the dose administered, their anticoagulant effect has a latency of onset of 1-3 days
C) Their dose is adjusted by repeated measurement of prothrombin time
D) They are contraindicated during pregnancy
6. Alkalinization of urine hastens the excretion of:
A) Weakly basic drugs B) Weakly acidic drugs
C) Strongly acidic drugs D) Nonpolar drugs
7. If the total amount of a drug present in the body at a given moment is 2.0 g and its plasma concentration is 50 µg/ml, its volume of distribution is:
A) 100 L B) 40 L C) 60 L D) 50 L
8. Induction of drug metabolizing enzymes involves:
A) A conformational change in the enzyme protein to favour binding of substrate molecules
B) Expression of enzyme molecules on the surface of hepatocytes
C) Enhanced transport of substrate molecules into hepatocytes
D) Increased synthesis of enzyme protein
9. Receptor agonists possess:
A) Affinity but no intrinsic activity
B) Intrinsic activity but no affinity
C) Affinity and intrinsic activity with a + sign
D) Affinity and intrinsic activity with a – sign

10. The choline ester resistant to both true and pseudocholinesterase is:
A) Methacholine B) Bethanechol C) Benzoylcholine D) Butyrylcholine
11. Sumatriptan acts by the following mechanism
A) 5HT 1D antagonist B) 5HT 1D/B agonist
C) 5HT 3 antagonist D) 5HT 1D/B antagonist
12. The principal process which terminates the action of noradrenaline released from adrenergic nerve ending is:
A) Degradation by MAO B) Methylation by COMT
C) Axonal uptake D) Extra neuronal uptake
13. Select the H1 antihistaminic which is used topically in the nose for allergic rhinitis:
A) Loratadine B) Cetirizine C) Fexofenadine D) Azelastine
14. Aspirin is contraindicated in pregnant women near term because:
A) Labour may be delayed and prolonged
B) Blood loss during delivery may be more
C) Foetus may suffer premature closure of ductus arteriosus
D) All of the above
15. Bronchodilators are useful in cough:
A) Only when cough is nonproductive
B) Only when cough is associated with thick sticky secretions
C) Only when reflex bronchoconstriction is associated
D) Irrespective of nature of cough or associated features
16. No surgical operation should be performed during the following stage of anaesthesia:
A) Stage I B) Stage II C) Stage III, plane 1 D) Stage III, plane 3
17. Benzodiazepines differ from barbiturates in the following aspects except:
A) They have a steeper dose response curve
B) They have higher therapeutic index
C) They have lower abuse liability
D) They do not induce microsomal drug metabolizing enzymes
18. The following drug combination should not be used in parkinsonism:
A) Levodopa + anticholinergics B) Levodopa + amantadine
C) Bromocriptine + levodopa D) Amantadine + anticholinergics
19. The following side effect of neuroleptics responds to propranolol:
A) Parkinsonism B) Acute muscle dystonia
C) Tardive dyskinesia D) Akathisia
20. Limitations of typical tricyclic anti-depressants include the following except:
A) Narrow safety margin B) Low oral bioavailability
C) Frequent side effects D) Long latent period for response

THE TAMIL NADU DR. M.G.R. MEDICAL UNIVERSITY

[MBBS 0325]

MARCH 2025

Sub. Code :6061

M.B.B.S. DEGREE EXAMINATIONS

(For the candidates admitted from the Academic Year 2019 - 2020)

SECOND PROFESSIONAL – SUPPLEMENTARY - (CBME)

PAPER I – PHARMACOLOGY

Q.P. Code: 526061

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - A

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Choose the drug with short plasma half life
A) Esmolol B) Chloroquine C) Digoxin D) Etanercept
2. Rapid development of Tolerance is
A) Resistance B) Teratogenicity C) Cumulation D) Tachyphylaxis
3. Nonequilibrium type alpha adrenergic blocker is
A) Ergotamine B) Phenoxybenzamine C) Phentolamine D) Tolazoline
4. Prolonged use of Hydralazine for the treatment of hypertension is likely to cause
A) Gynaecomastia B) Thrombocytopenia
C) Haemolytic Anemia D) Lupus Erythematosus
5. Name the beta blocker used in Congestive cardiac failure
A) Propranolol B) Metoprolol C) Atenolol D) Acebutolol
6. The prostaglandin analog used in Primary Pulmonary Hypertension
A) Carboprost B) Misoprostol C) Dinoprostone D) Epoprostenol
7. Propofol is the most preferred IV anaesthetic due to
A) Quick induction and rapid recovery B) Less respiratory depression
C) No change in blood pressure D) Positive inotropic effect
8. Selective COX-2 inhibitors are preferred as analgesic due to less risk of
A) Ischaemic heart disease B) Stroke C) Peptic ulcer D) Nephrotoxicity
9. Which among the following is NOT administered orally in bronchial asthma
A) Doxophylline B) Ketotifen C) Montelukast D) Sodium cromoglycate
10. Drug which precipitates acute intermittent porphyria is
A) Ketamine B) Thiopentone C) Halothane D) Propofol
11. An 'orphan drug' is:
A) A very cheap drug B) A drug which has no therapeutic use
C) A drug for treatment of a rare disease D) A drug which acts on Orphan receptors

12. Which of the following is a prodrug of adrenaline used topically in glaucoma:
A) Brimonidine B) Dipivefrine C) Apraclonidine D) Dorzolamide
13. Morphine stimulates:
A) Vasomotor centre B) Edinger Westphal nucleus
C) Temperature regulating centre D) Cough centre
14. Which of the following drug undergoes 'Hofmann' elimination:
A) Succinylcholine B) Pancuronium C) Vecuronium D) Atracurium
15. Organic nitrates relax vascular smooth muscle by:
A) Increasing intracellular cyclic GMP B) Increasing intracellular cyclic AMP
C) Decreasing intracellular cyclic AMP D) Decreasing intracellular cyclic GMP
16. Which of the following is an irreversible inhibitor of cyclooxygenase:
A) Aspirin B) Naproxen C) Indomethacin D) Piroxicam
17. Select the H₁ antihistaminic used topically for allergic rhinitis:
A) Loratadine B) Cetirizine C) Fexofenadine D) Azelastine
18. The following is a selective 5-HT_{1D} receptor agonist:
A) Buspirone B) Ondansetron C) Sumatriptan D) Ergotamine
19. Recombinant human erythropoietin is indicated for:
A) Megaloblastic anaemia B) Haemolytic anaemia
C) Anaemia in patients of thalassemia D) Anaemia in chronic renal failure
20. The characteristic adverse effect of HMG-CoA reductase inhibitors is:
A) Onycholysis B) Myopathy C) Alopecia D) Haemolysis

[MBBS 0325]

THE TAMIL NADU DR. M.G.R. MEDICAL UNIVERSITY

[MBBS 0325]

MARCH 2025

Sub. Code :6061

M.B.B.S. DEGREE EXAMINATIONS

(For the candidates admitted from the Academic Year 2019 - 2020)

SECOND PROFESSIONAL – SUPPLEMENTARY - (CBME)

PAPER I – PHARMACOLOGY

Q.P. Code: 526061

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - B

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Name the beta blocker used in Congestive cardiac failure
A) Propranolol B) Metoprolol C) Atenolol D) Acebutolol
2. The prostaglandin analog used in Primary Pulmonary Hypertension
A) Carboprost B) Misoprostol C) Dinoprostone D) Epoprostenol
3. Propofol is the most preferred IV anaesthetic due to
A) Quick induction and rapid recovery B) Less respiratory depression
C) No change in blood pressure D) Positive inotropic effect
4. Selective COX-2 inhibitors are preferred as analgesic due to less risk of
A) Ischaemic heart disease B) Stroke C) Peptic ulcer D) Nephrotoxicity
5. Which among the following is NOT administered orally in bronchial asthma
A) Doxophylline B) Ketotifen C) Montelukast D) Sodium cromoglycate
6. Drug which precipitates acute intermittent porphyria is
A) Ketamine B) Thiopentone C) Halothane D) Propofol
7. An 'orphan drug' is:
A) A very cheap drug B) A drug which has no therapeutic use
C) A drug for treatment of a rare disease D) A drug which acts on Orphan receptors
8. Which of the following is a prodrug of adrenaline used topically in glaucoma:
A) Brimonidine B) Dipivefrine C) Apraclonidine D) Dorzolamide
9. Morphine stimulates:
A) Vasomotor centre B) Edinger Westphal nucleus
C) Temperature regulating centre D) Cough centre
10. Which of the following drug undergoes 'Hofmann' elimination:
A) Succinylcholine B) Pancuronium C) Vecuronium D) Atracurium

11. Organic nitrates relax vascular smooth muscle by:
A) Increasing intracellular cyclic GMP B) Increasing intracellular cyclic AMP
C) Decreasing intracellular cyclic AMP D) Decreasing intracellular cyclic GMP
12. Which of the following is an irreversible inhibitor of cyclooxygenase:
A) Aspirin B) Naproxen C) Indomethacin D) Piroxicam
13. Select the H₁ antihistaminic used topically for allergic rhinitis:
A) Loratadine B) Cetirizine C) Fexofenadine D) Azelastine
14. The following is a selective 5-HT_{1D} receptor agonist:
A) Buspirone B) Ondansetron C) Sumatriptan D) Ergotamine
15. Recombinant human erythropoietin is indicated for:
A) Megaloblastic anaemia B) Haemolytic anaemia
C) Anaemia in patients of thalassemia D) Anaemia in chronic renal failure
16. The characteristic adverse effect of HMG-CoA reductase inhibitors is:
A) Onycholysis B) Myopathy C) Alopecia D) Haemolysis
17. Choose the drug with short plasma half life
A) Esmolol B) Chloroquine C) Digoxin D) Etanercept
18. Rapid development of Tolerance is
A) Resistance B) Teratogenicity C) Cumulation D) Tachyphylaxis
19. Nonequilibrium type alpha adrenergic blocker is
A) Ergotamine B) Phenoxybenzamine C) Phentolamine D) Tolazoline
20. Prolonged use of Hydralazine for the treatment of hypertension is likely to cause
A) Gynaecomastia B) Thrombocytopenia
C) Haemolytic Anemia D) Lupus Erythematosus

[MBBS 0325]

THE TAMIL NADU DR. M.G.R. MEDICAL UNIVERSITY

[MBBS 0325]

MARCH 2025

Sub. Code :6061

M.B.B.S. DEGREE EXAMINATIONS

(For the candidates admitted from the Academic Year 2019 - 2020)

SECOND PROFESSIONAL – SUPPLEMENTARY - (CBME)

PAPER I – PHARMACOLOGY

Q.P. Code: 526061

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - C

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Drug which precipitates acute intermittent porphyria is
A) Ketamine B) Thiopentone C) Halothane D) Propofol
2. An 'orphan drug' is:
A) A very cheap drug B) A drug which has no therapeutic use
C) A drug for treatment of a rare disease D) A drug which acts on Orphan receptors
3. Which of the following is a prodrug of adrenaline used topically in glaucoma:
A) Brimonidine B) Dipivefrine C) Apraclonidine D) Dorzolamide
4. Morphine stimulates:
A) Vasomotor centre B) Edinger Westphal nucleus
C) Temperature regulating centre D) Cough centre
5. Which of the following drug undergoes 'Hofmann' elimination:
A) Succinylcholine B) Pancuronium C) Vecuronium D) Atracurium
6. Organic nitrates relax vascular smooth muscle by:
A) Increasing intracellular cyclic GMP B) Increasing intracellular cyclic AMP
C) Decreasing intracellular cyclic AMP D) Decreasing intracellular cyclic GMP
7. Which of the following is an irreversible inhibitor of cyclooxygenase:
A) Aspirin B) Naproxen C) Indomethacin D) Piroxicam
8. Select the H₁ antihistaminic used topically for allergic rhinitis:
A) Loratadine B) Cetirizine C) Fexofenadine D) Azelastine
9. The following is a selective 5-HT_{1D} receptor agonist:
A) Buspirone B) Ondansetron C) Sumatriptan D) Ergotamine
10. Recombinant human erythropoietin is indicated for:
A) Megaloblastic anaemia B) Haemolytic anaemia
C) Anaemia in patients of thalassemia D) Anaemia in chronic renal failure

11. The characteristic adverse effect of HMG-CoA reductase inhibitors is:
A) Onycholysis B) Myopathy C) Alopecia D) Haemolysis
12. Choose the drug with short plasma half life
A) Esmolol B) Chloroquine C) Digoxin D) Etanercept
13. Rapid development of Tolerance is
A) Resistance B) Teratogenicity C) Cumulation D) Tachyphylaxis
14. Nonequilibrium type alpha adrenergic blocker is
A) Ergotamine B) Phenoxybenzamine C) Phentolamine D) Tolazoline
15. Prolonged use of Hydralazine for the treatment of hypertension is likely to cause
A) Gynaecomastia B) Thrombocytopenia
C) Haemolytic Anemia D) Lupus Erythematosus
16. Name the beta blocker used in Congestive cardiac failure
A) Propranolol B) Metoprolol C) Atenolol D) Acebutolol
17. The prostaglandin analog used in Primary Pulmonary Hypertension
A) Carboprost B) Misoprostol C) Dinoprostone D) Epoprostenol
18. Propofol is the most preferred IV anaesthetic due to
A) Quick induction and rapid recovery B) Less respiratory depression
C) No change in blood pressure D) Positive inotropic effect
19. Selective COX-2 inhibitors are preferred as analgesic due to less risk of
A) Ischaemic heart disease B) Stroke C) Peptic ulcer D) Nephrotoxicity
20. Which among the following is NOT administered orally in bronchial asthma
A) Doxophylline B) Ketotifen C) Montelukast D) Sodium cromoglycate

[MBBS 0325]

THE TAMIL NADU DR. M.G.R. MEDICAL UNIVERSITY

[MBBS 0325]

MARCH 2025

Sub. Code :6061

M.B.B.S. DEGREE EXAMINATIONS

(For the candidates admitted from the Academic Year 2019 - 2020)

SECOND PROFESSIONAL – SUPPLEMENTARY - (CBME)

PAPER I – PHARMACOLOGY

Q.P. Code: 526061

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - D

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Organic nitrates relax vascular smooth muscle by:
A) Increasing intracellular cyclic GMP B) Increasing intracellular cyclic AMP
C) Decreasing intracellular cyclic AMP D) Decreasing intracellular cyclic GMP
2. Which of the following is an irreversible inhibitor of cyclooxygenase:
A) Aspirin B) Naproxen C) Indomethacin D) Piroxicam
3. Select the H₁ antihistaminic used topically for allergic rhinitis:
A) Loratadine B) Cetirizine C) Fexofenadine D) Azelastine
4. The following is a selective 5-HT_{1D} receptor agonist:
A) Buspirone B) Ondansetron C) Sumatriptan D) Ergotamine
5. Recombinant human erythropoietin is indicated for:
A) Megaloblastic anaemia B) Haemolytic anaemia
C) Anaemia in patients of thalassemia D) Anaemia in chronic renal failure
6. The characteristic adverse effect of HMG-CoA reductase inhibitors is:
A) Onycholysis B) Myopathy C) Alopecia D) Haemolysis
7. Choose the drug with short plasma half life
A) Esmolol B) Chloroquine C) Digoxin D) Etanercept
8. Rapid development of Tolerance is
A) Resistance B) Teratogenicity C) Cumulation D) Tachyphylaxis
9. Nonequilibrium type alpha adrenergic blocker is
A) Ergotamine B) Phenoxybenzamine C) Phentolamine D) Tolazoline
10. Prolonged use of Hydralazine for the treatment of hypertension is likely to cause
A) Gynaecomastia B) Thrombocytopenia
C) Haemolytic Anemia D) Lupus Erythematosus

11. Name the beta blocker used in Congestive cardiac failure
A) Propranolol B) Metoprolol C) Atenolol D) Acebutolol
12. The prostaglandin analog used in Primary Pulmonary Hypertension
A) Carboprost B) Misoprostol C) Dinoprostone D) Epoprostenol
13. Propofol is the most preferred IV anaesthetic due to
A) Quick induction and rapid recovery B) Less respiratory depression
C) No change in blood pressure D) Positive inotropic effect
14. Selective COX-2 inhibitors are preferred as analgesic due to less risk of
A) Ischaemic heart disease B) Stroke C) Peptic ulcer D) Nephrotoxicity
15. Which among the following is NOT administered orally in bronchial asthma
A) Doxophylline B) Ketotifen C) Montelukast D) Sodium cromoglycate
16. Drug which precipitates acute intermittent porphyria is
A) Ketamine B) Thiopentone C) Halothane D) Propofol
17. An 'orphan drug' is:
A) A very cheap drug B) A drug which has no therapeutic use
C) A drug for treatment of a rare disease D) A drug which acts on Orphan receptors
18. Which of the following is a prodrug of adrenaline used topically in glaucoma:
A) Brimonidine B) Dipivefrine C) Apraclonidine D) Dorzolamide
19. Morphine stimulates:
A) Vasomotor centre B) Edinger Westphal nucleus
C) Temperature regulating centre D) Cough centre
20. Which of the following drug undergoes 'Hofmann' elimination:
A) Succinylcholine B) Pancuronium C) Vecuronium D) Atracurium

[MBBS 0325]

M.B.B.S. DEGREE EXAMINATION
(For the candidates admitted from the Academic Year 2019 - 2020)

SECOND PROFESSIONAL – (CBME)
PAPER I – PHARMACOLOGY

Q.P. Code: 526061

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - A

III. Multiple Choice Questions:

(20 x 1 = 20)

1. High plasma protein binding:
 - A) Increases volume of distribution of the drug
 - B) Facilitates glomerular filtration of the drug
 - C) Minimises drug interactions
 - D) Decreases volume of distribution of the drug
2. Microsomal enzyme induction can be a cause of:
 - A) Tolerance
 - B) Physical dependence
 - C) Psychological dependence
 - D) Placebo effect
3. The plasma half-life of penicillin-G is longer in the new born because their:
 - A) Plasma protein level is low
 - B) Drug metabolizing enzymes are immature
 - C) Glomerular filtration rate is low
 - D) Tubular transport mechanisms are not well developed
4. Receptors perform the following function/functions:
 - A) Ligand recognition
 - B) Signal transduction
 - C) Both ligand recognition and signal transduction
 - D) Disposal of agonists and antagonists
5. Which of the following is a relatively cerebro selective anticholinesterase found to afford symptomatic improvement in Alzheimer's disease:
 - A) Donepezil
 - B) Gemfibrozil
 - C) Pyridostigmine
 - D) Pyritinol
6. Hyoscine differs from atropine in that it:
 - A) Exerts depressant effects on the CNS at relatively low doses
 - B) Exerts more potent effects on the heart than on the eye
 - C) Is longer acting
 - D) Has weaker anti motion sickness activity
7. Choose the drug which is used as a short-term inotropic in severe congestive heart failure and has selective adrenergic β_1 agonistic activity but no dopaminergic agonistic activity:
 - A) Dopamine
 - B) Dobutamine
 - C) Amrinone
 - D) Salmeterol
8. Select the ergot compound which is primarily used for dementia:
 - A) Bromocriptine
 - B) Ergotamine
 - C) Codergocrine
 - D) Methysergide

9. Sulfasalazine is used in the following disease(s):
A) Bacillary dysentery
B) Ulcerative colitis
C) Rheumatoid arthritis
D) Both 'B' and 'C' are correct
10. Reflex bronchoconstriction is most likely to occur with the following form of inhaled anti-asthma medication:
A) Metered dose spray of drug in solution
B) Dry powder rota cap
C) Nebuliser
D) Nebuliser with spacer
11. Select the anaesthetic that increases the blood pressure:
A) Halothane
B) Fentanyl
C) Ketamine
D) Diazepam
12. Select the anti-convulsant drug that acts as a GABA transaminase inhibitor:
A) Gabapentin
B) Vigabatrin
C) Lamotrigine
D) Clobazam
13. The major limitation in the use of clozapine for treatment of schizophrenia is:
A) Its potential to cause agranulocytosis
B) Its inability to benefit negative symptoms of schizophrenia
C) High incidence of extrapyramidal side effects
D) Production of hyperprolactinemia
14. Choose the correct statement about the use of propranolol in anxiety:
A) Being non-sedative, it is the drug of choice in chronic anxiety states
B) It suppresses the psychological component of anxiety
C) It suppresses the autonomic manifestations of acutely stressful situations
D) Both 'B' and 'C' are correct
15. Strychnine produces convulsions by:
A) Stimulating NMDA receptors
B) Facilitating the excitatory transmitter glutamate
C) Blocking the inhibitory transmitter GABA
D) Blocking the inhibitory transmitter glycine
16. Beneficial effect / effects of diuretics in CHF patients include the following:
A) Symptomatic relief
B) Regression of pathological changes
C) Prolongation of life expectancy
D) Both 'A' and 'C'
17. Organic nitrates have predominantly venodilator action because:
A) They are selectively concentrated in veins
B) Veins express larger quantities of enzymes that generate nitric oxide from nitrates
C) Venous smooth muscle has greater capacity to relax
D) All of the above are correct
18. Coronary steal phenomenon has been noted most frequently with:
A) Glyceryl trinitrate
B) Dipyridamole
C) Propranolol
D) Diltiazem
19. Recombinant human erythropoietin is indicated for:
A) Megaloblastic anaemia
B) Haemolytic anaemia
C) Patients of thalassemia
D) Anaemia in chronic renal failure patients
20. Which of the following hypolipidemic drugs is most effective in raising HDL-cholesterol level?
A) Nicotinic acid
B) Fenofibrate
C) Cholestyramine
D) Pravastatin

M.B.B.S. DEGREE EXAMINATION
(For the candidates admitted from the Academic Year 2019 - 2020)

SECOND PROFESSIONAL – (CBME)
PAPER I – PHARMACOLOGY

Q.P. Code: 526061

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - B

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Receptors perform the following function/functions:
A) Ligand recognition
B) Signal transduction
C) Both ligand recognition and signal transduction
D) Disposal of agonists and antagonists
2. Which of the following is a relatively cerebro selective anticholinesterase found to afford symptomatic improvement in Alzheimer's disease:
A) Donepezil B) Gemfibrozil C) Pyridostigmine D) Pyritinol
3. Hyoscine differs from atropine in that it:
A) Exerts depressant effects on the CNS at relatively low doses
B) Exerts more potent effects on the heart than on the eye
C) Is longer acting
D) Has weaker anti motion sickness activity
4. Choose the drug which is used as a short-term inotropic in severe congestive heart failure and has selective adrenergic β_1 agonistic activity but no dopaminergic agonistic activity:
A) Dopamine B) Dobutamine C) Amrinone D) Salmeterol
5. Select the ergot compound which is primarily used for dementia:
A) Bromocriptine B) Ergotamine C) Codergocrine D) Methysergide
6. Sulfasalazine is used in the following disease(s):
A) Bacillary dysentery B) Ulcerative colitis
C) Rheumatoid arthritis D) Both 'B' and 'C' are correct
7. Reflex bronchoconstriction is most likely to occur with the following form of inhaled anti-asthma medication:
A) Metered dose spray of drug in solution B) Dry powder rota cap
C) Nebuliser D) Nebuliser with spacer
8. Select the anaesthetic that increases the blood pressure:
A) Halothane B) Fentanyl C) Ketamine D) Diazepam
9. Select the anti-convulsant drug that acts as a GABA transaminase inhibitor:
A) Gabapentin B) Vigabatrin C) Lamotrigine D) Clobazam

10. The major limitation in the use of clozapine for treatment of schizophrenia is:
 - A) Its potential to cause agranulocytosis
 - B) Its inability to benefit negative symptoms of schizophrenia
 - C) High incidence of extrapyramidal side effects
 - D) Production of hyperprolactinemia
11. Choose the correct statement about the use of propranolol in anxiety:
 - A) Being non-sedative, it is the drug of choice in chronic anxiety states
 - B) It suppresses the psychological component of anxiety
 - C) It suppresses the autonomic manifestations of acutely stressful situations
 - D) Both 'B' and 'C' are correct
12. Strychnine produces convulsions by:
 - A) Stimulating NMDA receptors
 - B) Facilitating the excitatory transmitter glutamate
 - C) Blocking the inhibitory transmitter GABA
 - D) Blocking the inhibitory transmitter glycine
13. Beneficial effect / effects of diuretics in CHF patients include the following:
 - A) Symptomatic relief
 - B) Regression of pathological changes
 - C) Prolongation of life expectancy
 - D) Both 'A' and 'C'
14. Organic nitrates have predominantly venodilator action because:
 - A) They are selectively concentrated in veins
 - B) Veins express larger quantities of enzymes that generate nitric oxide from nitrates
 - C) Venous smooth muscle has greater capacity to relax
 - D) All of the above are correct
15. Coronary steal phenomenon has been noted most frequently with:
 - A) Glyceryl trinitrate
 - B) Dipyridamole
 - C) Propranolol
 - D) Diltiazem
16. Recombinant human erythropoietin is indicated for:
 - A) Megaloblastic anaemia
 - B) Haemolytic anaemia
 - C) Patients of thalassemia
 - D) Anaemia in chronic renal failure patients
17. Which of the following hypolipidemic drugs is most effective in raising HDL-cholesterol level?
 - A) Nicotinic acid
 - B) Fenofibrate
 - C) Cholestyramine
 - D) Pravastatin
18. High plasma protein binding:
 - A) Increases volume of distribution of the drug
 - B) Facilitates glomerular filtration of the drug
 - C) Minimises drug interactions
 - D) Decreases volume of distribution of the drug
19. Microsomal enzyme induction can be a cause of:
 - A) Tolerance
 - B) Physical dependence
 - C) Psychological dependence
 - D) Placebo effect
20. The plasma half-life of penicillin-G is longer in the new born because their:
 - A) Plasma protein level is low
 - B) Drug metabolizing enzymes are immature
 - C) Glomerular filtration rate is low
 - D) Tubular transport mechanisms are not well developed

M.B.B.S. DEGREE EXAMINATION
(For the candidates admitted from the Academic Year 2019 - 2020)

SECOND PROFESSIONAL – (CBME)
PAPER I – PHARMACOLOGY

Q.P. Code: 526061

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - C

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Select the ergot compound which is primarily used for dementia:
A) Bromocriptine B) Ergotamine C) Codergocrine D) Methysergide
2. Sulfasalazine is used in the following disease(s):
A) Bacillary dysentery B) Ulcerative colitis
C) Rheumatoid arthritis D) Both 'B' and 'C' are correct
3. Reflex bronchoconstriction is most likely to occur with the following form of inhaled anti-asthma medication:
A) Metered dose spray of drug in solution B) Dry powder rota cap
C) Nebuliser D) Nebuliser with spacer
4. Select the anaesthetic that increases the blood pressure:
A) Halothane B) Fentanyl C) Ketamine D) Diazepam
5. Select the anti-convulsant drug that acts as a GABA transaminase inhibitor:
A) Gabapentin B) Vigabatrin C) Lamotrigine D) Clobazam
6. The major limitation in the use of clozapine for treatment of schizophrenia is:
A) Its potential to cause agranulocytosis
B) Its inability to benefit negative symptoms of schizophrenia
C) High incidence of extrapyramidal side effects
D) Production of hyperprolactinemia
7. Choose the correct statement about the use of propranolol in anxiety:
A) Being non-sedative, it is the drug of choice in chronic anxiety states
B) It suppresses the psychological component of anxiety
C) It suppresses the autonomic manifestations of acutely stressful situations
D) Both 'B' and 'C' are correct
8. Strychnine produces convulsions by:
A) Stimulating NMDA receptors
B) Facilitating the excitatory transmitter glutamate
C) Blocking the inhibitory transmitter GABA
D) Blocking the inhibitory transmitter glycine
9. Beneficial effect / effects of diuretics in CHF patients include the following:
A) Symptomatic relief B) Regression of pathological changes
C) Prolongation of life expectancy D) Both 'A' and 'C'

10. Organic nitrates have predominantly venodilator action because:
 - A) They are selectively concentrated in veins
 - B) Veins express larger quantities of enzymes that generate nitric oxide from nitrates
 - C) Venous smooth muscle has greater capacity to relax
 - D) All of the above are correct
11. Coronary steal phenomenon has been noted most frequently with:
 - A) Glyceryl trinitrate B) Dipyridamole C) Propranolol D) Diltiazem
12. Recombinant human erythropoietin is indicated for:
 - A) Megaloblastic anaemia B) Haemolytic anaemia
 - C) Patients of thalassemia D) Anaemia in chronic renal failure patients
13. Which of the following hypolipidemic drugs is most effective in raising HDL-cholesterol level?
 - A) Nicotinic acid B) Fenofibrate C) Cholestyramine D) Pravastatin
14. High plasma protein binding:
 - A) Increases volume of distribution of the drug
 - B) Facilitates glomerular filtration of the drug
 - C) Minimises drug interactions
 - D) Decreases volume of distribution of the drug
15. Microsomal enzyme induction can be a cause of:
 - A) Tolerance B) Physical dependence
 - C) Psychological dependence D) Placebo effect
16. The plasma half-life of penicillin-G is longer in the new born because their:
 - A) Plasma protein level is low
 - B) Drug metabolizing enzymes are immature
 - C) Glomerular filtration rate is low
 - D) Tubular transport mechanisms are not well developed
17. Receptors perform the following function/functions:
 - A) Ligand recognition
 - B) Signal transduction
 - C) Both ligand recognition and signal transduction
 - D) Disposal of agonists and antagonists
18. Which of the following is a relatively cerebro selective anticholinesterase found to afford symptomatic improvement in Alzheimer's disease:
 - A) Donepezil B) Gemfibrozil C) Pyridostigmine D) Pyritinol
19. Hyoscine differs from atropine in that it:
 - A) Exerts depressant effects on the CNS at relatively low doses
 - B) Exerts more potent effects on the heart than on the eye
 - C) Is longer acting
 - D) Has weaker anti motion sickness activity
20. Choose the drug which is used as a short-term inotropic in severe congestive heart failure and has selective adrenergic β_1 agonistic activity but no dopaminergic agonistic activity:
 - A) Dopamine B) Dobutamine C) Amrinone D) Salmeterol

M.B.B.S. DEGREE EXAMINATION
(For the candidates admitted from the Academic Year 2019 - 2020)

SECOND PROFESSIONAL – (CBME)
PAPER I – PHARMACOLOGY

Q.P. Code: 526061

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - D

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Select the anti-convulsant drug that acts as a GABA transaminase inhibitor:
A) Gabapentin B) Vigabatrin C) Lamotrigine D) Clobazam
2. The major limitation in the use of clozapine for treatment of schizophrenia is:
A) Its potential to cause agranulocytosis
B) Its inability to benefit negative symptoms of schizophrenia
C) High incidence of extrapyramidal side effects
D) Production of hyperprolactinemia
3. Choose the correct statement about the use of propranolol in anxiety:
A) Being non-sedative, it is the drug of choice in chronic anxiety states
B) It suppresses the psychological component of anxiety
C) It suppresses the autonomic manifestations of acutely stressful situations
D) Both 'B' and 'C' are correct
4. Strychnine produces convulsions by:
A) Stimulating NMDA receptors
B) Facilitating the excitatory transmitter glutamate
C) Blocking the inhibitory transmitter GABA
D) Blocking the inhibitory transmitter glycine
5. Beneficial effect / effects of diuretics in CHF patients include the following:
A) Symptomatic relief B) Regression of pathological changes
C) Prolongation of life expectancy D) Both 'A' and 'C'
6. Organic nitrates have predominantly venodilator action because:
A) They are selectively concentrated in veins
B) Veins express larger quantities of enzymes that generate nitric oxide from nitrates
C) Venous smooth muscle has greater capacity to relax
D) All of the above are correct
7. Coronary steal phenomenon has been noted most frequently with:
A) Glyceryl trinitrate B) Dipyridamole C) Propranolol D) Diltiazem
8. Recombinant human erythropoietin is indicated for:
A) Megaloblastic anaemia B) Haemolytic anaemia
C) Patients of thalassemia D) Anaemia in chronic renal failure patients

9. Which of the following hypolipidemic drugs is most effective in raising HDL-cholesterol level?
A) Nicotinic acid B) Fenofibrate C) Cholestyramine D) Pravastatin
10. High plasma protein binding:
A) Increases volume of distribution of the drug
B) Facilitates glomerular filtration of the drug
C) Minimises drug interactions
D) Decreases volume of distribution of the drug
11. Microsomal enzyme induction can be a cause of:
A) Tolerance B) Physical dependence
C) Psychological dependence D) Placebo effect
12. The plasma half-life of penicillin-G is longer in the new born because their:
A) Plasma protein level is low
B) Drug metabolizing enzymes are immature
C) Glomerular filtration rate is low
D) Tubular transport mechanisms are not well developed
13. Receptors perform the following function/functions:
A) Ligand recognition
B) Signal transduction
C) Both ligand recognition and signal transduction
D) Disposal of agonists and antagonists
14. Which of the following is a relatively cerebro selective anticholinesterase found to afford symptomatic improvement in Alzheimer's disease:
A) Donepezil B) Gemfibrozil C) Pyridostigmine D) Pyritinol
15. Hyoscine differs from atropine in that it:
A) Exerts depressant effects on the CNS at relatively low doses
B) Exerts more potent effects on the heart than on the eye
C) Is longer acting
D) Has weaker anti motion sickness activity
16. Choose the drug which is used as a short-term inotropic in severe congestive heart failure and has selective adrenergic β_1 agonistic activity but no dopaminergic agonistic activity:
A) Dopamine B) Dobutamine C) Amrinone D) Salmeterol
17. Select the ergot compound which is primarily used for dementia:
A) Bromocriptine B) Ergotamine C) Codergocrine D) Methysergide
18. Sulfasalazine is used in the following disease(s):
A) Bacillary dysentery B) Ulcerative colitis
C) Rheumatoid arthritis D) Both 'B' and 'C' are correct
19. Reflex bronchoconstriction is most likely to occur with the following form of inhaled anti-asthma medication:
A) Metered dose spray of drug in solution B) Dry powder rota cap
C) Nebuliser D) Nebuliser with spacer
20. Select the anaesthetic that increases the blood pressure:
A) Halothane B) Fentanyl C) Ketamine D) Diazepam

M.B.B.S. DEGREE EXAMINATION
(For the candidates admitted from the Academic Year 2019 - 2020)

SECOND PROFESSIONAL – SUPPLEMENTARY – (CBME)
PAPER I – PHARMACOLOGY

Q.P. Code: 526061

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET – A

III. Multiple Choice Questions:

(20 x 1 = 20)

1. The Vasodilator peptide is
A) Endothelins
B) Calcitonin gene related peptide
C) Neuropeptide y
D) Bradykinin
2. Which one of the following is not a statement for rectal route of administration
A) Drugs are absorbed in internal haemorrhoidal vein
B) Irritant, unpleasant drugs can be given
C) It bypasses liver
D) Drugs like indomethacin, paracetamol can be given
3. Leflunomide acts by
A) Dihydro folate reductase inhibitor
B) Thiopurine methyl transferase inhibitor
C) Dihydro orotate dehydrogenase inhibitor
D) Cyclooxygenase inhibitor
4. Selective endothelin -A (ET-A) receptor antagonist is
A) Ambrisentan B) Bosentan C) Macitentan D) Atrial natriuretic peptide
5. Which one of the Atropine derivative is not used as mydriatic agent
A) Homatropine B) Cyclopentolate C) Tropicamide D) Pirenzepine
6. Ganglionic stimulant is
A) Hexamethonium B) Pentolinium C) Mecamylamine D) Varenicline
7. The selective alpha -2 agonist is
A) Phenylephrine B) Clonidine C) Alpha methyl dopa D) Ephedrine
8. Neprilysin inhibitor is
A) Sacubitril B) Icatibant C) Nesiritide D) Bosentan
9. The long-acting Nitrate is
A) Nitroglycerine B) Isosorbide dinitrate
C) Isosorbide-5-mono nitrate D) Pentaerythritol tetra nitrate

10. The parenteral anti-hypertensive drug therapy with short duration of action used in emergency is
A) Labetalol B) Clonidine C) Captopril D) Esmolol
11. Mannitol is indicated in
A) Acute Tubular necrosis B) Pulmonary edema
C) Acute left ventricular failure D) Acute congestive Glaucoma
12. Hofmann elimination is by
A) Disulfiram B) Atracurium C) Allopurinol D) Ciprofloxacin
13. In Active transport
A) Translocate the substrate in the direction of its electrochemical gradient
B) Transport solute against its electrochemical gradient
C) It does not require energy
D) Solute carrier transporters
14. Drug induced diseases called as
A) Dependence B) Tolerance
C) Physician induced diseases D) Tachyphylaxis
15. The Intrinsic sympathomimetic beta blocker is
A) Metoprolol B) Celiprolol C) Nebivolol D) Atenolol
16. Neostigmine antagonize the block of
A) d-Tubocurarine B) Succinyl choline C) Quinine D) Dantrolene sodium
17. The long-acting potent nonsteroidal anti-inflammatory drug is
A) Ketorolac B) Mephenamic acid C) Piroxicam D) Indomethacin
18. The inverse agonist of Benzodiazepines is
A) Muscimol B) Flumazenil C) Beta carboline D) Propofol
19. An inhibitor of GABA transaminase is
A) Tigabine B) Vigabatrin C) Gabapentin D) Topiramate
20. The Atypical Antipsychotic drug is
A) Levosulpiride B) Thioridazine C) Penfluridol D) Risperidone

[MBBS 1025]

M.B.B.S. DEGREE EXAMINATION
(For the candidates admitted from the Academic Year 2019 - 2020)

SECOND PROFESSIONAL – SUPPLEMENTARY – (CBME)
PAPER I – PHARMACOLOGY

Q.P. Code: 526061

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET – B

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Leflunomide acts by
 - A) Dihydro folate reductase inhibitor
 - B) Thiopurine methyl transferase inhibitor
 - C) Dihydro orotate dehydrogenase inhibitor
 - D) Cyclooxygenase inhibitor
2. Selective endothelin -A (ET-A) receptor antagonist is
 - A) Ambrisentan
 - B) Bosentan
 - C) Macitentan
 - D) Atrial natriuretic peptide
3. Which one of the Atropine derivative is not used as mydriatic agent
 - A) Homatropine
 - B) Cyclopentolate
 - C) Tropicamide
 - D) Pirenzepine
4. Ganglionic stimulant is
 - A) Hexamethonium
 - B) Pentolinium
 - C) Mecamylamine
 - D) Varenicline
5. The selective alpha -2 agonist is
 - A) Phenylephrine
 - B) Clonidine
 - C) Alpha methyl dopa
 - D) Ephedrine
6. Neprilysin inhibitor is
 - A) Sacubitril
 - B) Icatibant
 - C) Nesiritide
 - D) Bosentan
7. The long-acting Nitrate is
 - A) Nitroglycerine
 - B) Isosorbide dinitrate
 - C) Isosorbide-5-mono nitrate
 - D) Pentaerythritol tetra nitrate
8. The parenteral anti-hypertensive drug therapy with short duration of action used in emergency is
 - A) Labetalol
 - B) Clonidine
 - C) Captopril
 - D) Esmolol
9. Mannitol is indicated in
 - A) Acute Tubular necrosis
 - B) Pulmonary edema
 - C) Acute left ventricular failure
 - D) Acute congestive Glaucoma

10. Hofmann elimination is by
A) Disulfiram B) Atracurium C) Allopurinol D) Ciprofloxacin
11. In Active transport
A) Translocate the substrate in the direction of its electrochemical gradient
B) Transport solute against its electrochemical gradient
C) It does not require energy
D) Solute carrier transporters
12. Drug induced diseases called as
A) Dependence B) Tolerance
C) Physician induced diseases D) Tachyphylaxis
13. The Intrinsic sympathomimetic beta blocker is
A) Metoprolol B) Celiprolol C) Nebivolol D) Atenolol
14. Neostigmine antagonize the block of
A) d-Tubocurarine B) Succinyl choline C) Quinine D) Dantrolene sodium
15. The long-acting potent nonsteroidal anti-inflammatory drug is
A) Ketorolac B) Mephenamic acid C) Piroxicam D) Indomethacin
16. The inverse agonist of Benzodiazepines is
A) Muscimol B) Flumazenil C) Beta carboline D) Propofol
17. An inhibitor of GABA transaminase is
A) Tigabine B) Vigabatrin C) Gabapentin D) Topiramate
18. The Atypical Antipsychotic drug is
A) Levosulpiride B) Thioridazine C) Penfluridol D) Risperidone
19. The Vasodilator peptide is
A) Endothelins B) Calcitonin gene related peptide
C) Neuropeptide y D) Bradykinin
20. Which one of the following is not a statement for rectal route of administration
A) Drugs are absorbed in internal haemorrhoidal vein
B) Irritant, unpleasant drugs can be given
C) It bypasses liver
D) Drugs like indomethacin, paracetamol can be given

M.B.B.S. DEGREE EXAMINATION
(For the candidates admitted from the Academic Year 2019 - 2020)

SECOND PROFESSIONAL – SUPPLEMENTARY – (CBME)
PAPER I – PHARMACOLOGY

Q.P. Code: 526061

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET – C

III. Multiple Choice Questions:

(20 x 1 = 20)

1. The selective alpha -2 agonist is
A) Phenylephrine B) Clonidine C) Alpha methyl dopa D) Ephedrine
2. Neprilysin inhibitor is
A) Sacubitril B) Icatibant C) Nesiritide D) Bosentan
3. The long-acting Nitrate is
A) Nitroglycerine B) Isosorbide dinitrate
C) Isosorbide-5-mono nitrate D) Pentaerythritol tetra nitrate
4. The parenteral anti-hypertensive drug therapy with short duration of action used in emergency is
A) Labetalol B) Clonidine C) Captopril D) Esmolol
5. Mannitol is indicated in
A) Acute Tubular necrosis B) Pulmonary edema
C) Acute left ventricular failure D) Acute congestive Glaucoma
6. Hofmann elimination is by
A) Disulfiram B) Atracurium C) Allopurinol D) Ciprofloxacin
7. In Active transport
A) Translocate the substrate in the direction of its electrochemical gradient
B) Transport solute against its electrochemical gradient
C) It does not require energy
D) Solute carrier transporters
8. Drug induced diseases called as
A) Dependence B) Tolerance
C) Physician induced diseases D) Tachyphylaxis
9. The Intrinsic sympathomimetic beta blocker is
A) Metoprolol B) Celiprolol C) Nebivolol D) Atenolol

10. Neostigmine antagonize the block of
A) d-Tubocurarine B) Succinyl choline C) Quinine D) Dantrolene sodium
11. The long-acting potent nonsteroidal anti-inflammatory drug is
A) Ketorolac B) Mephenamic acid C) Piroxicam D) Indomethacin
12. The inverse agonist of Benzodiazepines is
A) Muscimol B) Flumazenil C) Beta carboline D) Propofol
13. An inhibitor of GABA transaminase is
A) Tigabine B) Vigabatrin C) Gabapentin D) Topiramate
14. The Atypical Antipsychotic drug is
A) Levosulpiride B) Thioridazine C) Penfluridol D) Risperidone
15. The Vasodilator peptide is
A) Endothelins B) Calcitonin gene related peptide
C) Neuropeptide y D) Bradykinin
16. Which one of the following is not a statement for rectal route of administration
A) Drugs are absorbed in internal haemorrhoidal vein
B) Irritant, unpleasant drugs can be given
C) It bypasses liver
D) Drugs like indomethacin, paracetamol can be given
17. Leflunomide acts by
A) Dihydro folate reductase inhibitor
B) Thiopurine methyl transferase inhibitor
C) Dihydro orotate dehydrogenase inhibitor
D) Cyclooxygenase inhibitor
18. Selective endothelin -A (ET-A) receptor antagonist is
A) Ambrisentan B) Bosentan C) Macitentan D) Atrial natriuretic peptide
19. Which one of the Atropine derivative is not used as mydriatic agent
A) Homatropine B) Cyclopentolate C) Tropicamide D) Pirenzepine
20. Ganglionic stimulant is
A) Hexamethonium B) Pentolinium C) Mecamylamine D) Varenicline

[MBBS 1025]

M.B.B.S. DEGREE EXAMINATION
(For the candidates admitted from the Academic Year 2019 - 2020)

SECOND PROFESSIONAL – SUPPLEMENTARY – (CBME)
PAPER I – PHARMACOLOGY

Q.P. Code: 526061

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET – D

III. Multiple Choice Questions:

(20 x 1 = 20)

1. The parenteral anti-hypertensive drug therapy with short duration of action used in emergency is
A) Labetalol B) Clonidine C) Captopril D) Esmolol
2. Mannitol is indicated in
A) Acute Tubular necrosis B) Pulmonary edema
C) Acute left ventricular failure D) Acute congestive Glaucoma
3. Hofmann elimination is by
A) Disulfiram B) Atracurium C) Allopurinol D) Ciprofloxacin
4. In Active transport
A) Translocate the substrate in the direction of its electrochemical gradient
B) Transport solute against its electrochemical gradient
C) It does not require energy
D) Solute carrier transporters
5. Drug induced diseases called as
A) Dependence B) Tolerance
C) Physician induced diseases D) Tachyphylaxis
6. The Intrinsic sympathomimetic beta blocker is
A) Metoprolol B) Celiprolol C) Nebivolol D) Atenolol
7. Neostigmine antagonize the block of
A) d-Tubocurarine B) Succinyl choline C) Quinine D) Dantrolene sodium
8. The long-acting potent nonsteroidal anti-inflammatory drug is
A) Ketorolac B) Mephenamic acid C) Piroxicam D) Indomethacin
9. The inverse agonist of Benzodiazepines is
A) Muscimol B) Flumazenil C) Beta carboline D) Propofol

10. An inhibitor of GABA transaminase is
A) Tigabine B) Vigabatrin C) Gabapentin D) Topiramate
11. The Atypical Antipsychotic drug is
A) Levosulpiride B) Thioridazine C) Penfluridol D) Risperidone
12. The Vasodilator peptide is
A) Endothelins B) Calcitonin gene related peptide
C) Neuropeptide y D) Bradykinin
13. Which one of the following is not a statement for rectal route of administration
A) Drugs are absorbed in internal haemorrhoidal vein
B) Irritant, unpleasant drugs can be given
C) It bypasses liver
D) Drugs like indomethacin, paracetamol can be given
14. Leflunomide acts by
A) Dihydro folate reductase inhibitor
B) Thiopurine methyl transferase inhibitor
C) Dihydro orotate dehydrogenase inhibitor
D) Cyclooxygenase inhibitor
15. Selective endothelin -A (ET-A) receptor antagonist is
A) Ambrisentan B) Bosentan C) Macitentan D) Atrial natriuretic peptide
16. Which one of the Atropine derivative is not used as mydriatic agent
A) Homatropine B) Cyclopentolate C) Tropicamide D) Pirenzepine
17. Ganglionic stimulant is
A) Hexamethonium B) Pentolinium C) Mecamylamine D) Varenicline
18. The selective alpha -2 agonist is
A) Phenylephrine B) Clonidine C) Alpha methyl dopa D) Ephedrine
19. Neprilysin inhibitor is
A) Sacubitril B) Icatibant C) Nesiritide D) Bosentan
20. The long-acting Nitrate is
A) Nitroglycerine B) Isosorbide dinitrate
C) Isosorbide-5-mono nitrate D) Pentaerythritol tetra nitrate

[MBBS 1025]