

M.B.B.S. DEGREE EXAMINATION
(For the candidates admitted from the Academic Year 2019-2020)

THIRD PROFESSIONAL PART II - CBME
PAPER II – OBSTETRICS & GYNAECOLOGY

Q.P. Code: 526086

Time: 30 Minutes

Maximum : 20 Marks

Answer All Questions

Choose one correct answer in the box provided in the Answer Script.
No overwriting should be done. Choice should be given in Capital Letters.

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Day of ovulation is
 - A) 14 days before next menstruation
 - B) 14 days after menstruation
 - C) 10 days after menstruation
 - D) 18 days after menstruation
2. All of the following can occur due to DES exposure inutero in female baby except
 - A) T shaped uterus
 - B) Clear cell carcinoma
 - C) Vaginal adenosis
 - D) Renal anomalies
3. Signet ring cells in a cellular stroma is diagnostic of
 - A) Gonadoblastoma
 - B) Hilus cell tumour
 - C) Krukenberg tumour
 - D) Struma ovarii
4. A women with which of the following health problem must avoid centchroman
 - A) PCOS
 - B) Endometrial hyperplasia
 - C) DUB
 - D) Endometriosis
5. Prostate gland is homologous to
 - A) Bartholins gland
 - B) Cowper's gland
 - C) Skene glands
 - D) Bulbourethral gland
6. Tubal causes of infertility is diagnosed by all except
 - A) HSG
 - B) SSG
 - C) Lap chromopertubation
 - D) Post coital test
7. Surrogacy is indicated in all except
 - A) Premature ovarian failure
 - B) Absent uterus
 - C) Hereditary diseases
 - D) Elderly mother
8. HPV – which serotype is responsible for cancer cervix
 - A) 6
 - B) 11
 - C) 16
 - D) 32
9. All are true regarding endometriosis except
 - A) Hormone dependent
 - B) Contains clear fluid
 - C) Involve lung and pleura
 - D) Ovary is the most common site
10. Halban's sign is seen in
 - A) Endometriosis
 - B) Fibroid uterus
 - C) Adenomyosis
 - D) Ca cervix

11. Diagnostic of menopause
A) Serum FSH less than 40 B) Serum FSH more than 40
C) Serum LH more than 20 D) Serum Estradiol less than 30
12. Most common cause of hirsutism in a young female
A) Ovarian tumour B) Adrenal tumour
C) Congenital adrenal hyperplasia D) PCOS
13. Difficulties in Minimally Invasive Surgery are all except
A) Indirect palpation of tissues B) Restricted movement of tools
C) Presence of three dimensional vision D) Counter- intuitive motion
14. Vertical fusion defect can result in
A) Septate uterus B) Transverse vaginal septum
C) Uterine didelphus D) Bicornuate uterus
15. Ulceration of the vulva is seen in all except
A) Bacterial vaginosis B) Syphilis C) Chancroid D) Behcet's disease
16. Not a barrier contraceptive
A) Diaphragm B) centchroman C) Condom D) Today
17. Ureter is identified intraoperatively by
A) Rich arterial plexus B) Relation to lumbar plexus
C) Peristaltic movement D) Close to renal vein
18. Cause of decubitus ulcer in prolapse
A) Friction B) Venous congestion C) Intercourse D) Trauma
19. Emergency contraception prevents pregnancy by all except
A) Delaying ovulation B) Inhibiting fertilisation
C) Preventing implantation D) Interrupting early pregnancy
20. Most common gene mutation in fibroid
A) MED-12 B) COL 4 A5-A6 C) FH D) None of the above

[MBBS 0324]

M.B.B.S. DEGREE EXAMINATION

(For the candidates admitted from the Academic Year 2019-2020)

THIRD PROFESSIONAL PART II - SUPPLEMENTARY (CBME)

PAPER II – OBSTETRICS & GYNAECOLOGY

Q.P. Code: 526086

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

**Choose one correct answer in the box provided in the Answer Script.
No overwriting should be done. Choice should be given in Capital Letters.**

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Chemotherapy of choriocarcinoma is with
A) Methotrexate B) Bleomycin C) Busulphan D) Cyclophosphamide
2. Commonest malignancy of body of the uterus is
A) Adenoacanthoma B) Sarcoma
C) Squamous cell carcinoma D) Adenocarcinoma
3. The spinbarkeit of cervical mucous is maximum in
A) Early proliferative phase B) Ovulatory phase
C) Secretory phase D) Menstrual phase
4. Most common site of genital tuberculosis
A) Fallopian tube B) Uterus C) Ovary D) Vagina
5. Commonest manifestation of endometriosis is
A) Infertility B) Pain C) Leukorrhoea D) Bleeding
6. The size of mature ovum
A) 100 - 110 μ B) 120 - 140 μ C) 140 - 160 μ D) 180 - 200 μ
7. For sex determination smear is taken from
A) Nose B) Buccal mucosa C) Vagina D) Skin
8. Strawberry vagina is seen in infection with
A) Hemophilus vaginalis B) Trichomonas vaginalis
C) Moniliasis D) Herpes simplex
9. MIRENA IUD contains
A) Levonorgestrel B) Regestrone
C) Medroxyprogesterone D) Ethinylestradiol
10. Treatment for nulliparous prolapse is
A) Manchesters operation B) Sling surgery
C) Vaginal hysterectomy D) Vaginal pessary

11. Tranexamic acid is useful in
A) Dyspareunia
B) Endometriosis
C) Menorrhagia
D) Stein Levinthal syndrome
12. Fertilisation occurs in
A) Ampulla
B) Isthmus
C) Fimbria
D) Uterus
13. Ovarian tumors causing masculinisation are
A) Arrhenoblastoma
B) Dysgerminoma
C) Gonadoblastoma
D) Granulosa cell tumor
14. Artificial insemination of husband semen is indicated in
A) Tubal factor
B) Cervical factor
C) Ovarian factor
D) Pituitary factor
15. Estrogens are contraindicated in
A) Senile vaginitis
B) Senile endometritis
C) Endometrial carcinoma
D) Contraception
16. Not a cause of hirsutism is
A) Granulosa cell tumor
B) Arrhenoblastoma
C) Polycystic ovary
D) Congenital adrenal hyperplasia
17. Secondary amenorrhoea after abortion is seen in
A) Anemia
B) Uterine atresia
C) Imperforate hymen
D) Asherman syndrome
18. Endometrial thickness in a postmenopausal women
A) 2 mm
B) 4 mm
C) 6 mm
D) 7 mm
19. In a lactating mother oral pills of choice is
A) Triphasic pill
B) Sequential pill
C) Mini pill
D) Combined pills
20. In young women deep conisation may lead to all except
A) Midtrimester abortion
B) Sterility
C) Premature labour
D) Cervical dystocia

[MBBS 0524]

11. The best diagnostic method of tubal block is
A) Ultrasonography B) Hysterosalpingography C) Colposcopy D) Hysteroscopy
12. The most common cause of hirsutism in a female is
A) Androblastoma B) Luteoma ovary
C) Adrenal hyperplasia D) PCOS
13. The following are common symptoms of endometriosis except
A) Diarrhoea B) Menorrhagia C) Dyspareunia D) Dysmenorrhoea
14. Gonococcal salpingitis is associated with all except
A) Pain B) Fever C) Vaginal bleeding D) Leukocytosis
15. The most common symptom of menopause
A) Insomnia B) Headache C) Hot flushes D) Depression
16. Nabothian follicles are seen in
A) Cervix B) Vagina C) Ovary D) Uterus
17. Commonest site of Myoma
A) Cervical B) Submucous C) Intramural D) Subserous
18. Which of the following organism causes condyloma acuminata
A) Fungus B) Virus C) Bacteria D) Protozoa
19. Granulosa and theca cell tumours are seen frequently
A) In infancy B) Before puberty C) In post-menopausal D) Reproductive age
20. The presence of pyometra in a post-menopausal woman is strongly suggestive of
A) Diabetes B) Tuberculous endometritis
C) Malignancy D) Degenerating myoma

[MBBS 0325]

THE TAMIL NADU DR. M.G.R. MEDICAL UNIVERSITY

[MBBS 0325]

MARCH 2025

Sub. Code :6086

**M.B.B.S. DEGREE EXAMINATION
(For the candidates admitted from the Academic Year 2019-2020)**

**THIRD PROFESSIONAL MBBS PART II
PAPER II – OBSTETRICS & GYNAECOLOGY
Q.P. Code: 526086**

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - B

III. Multiple Choice Questions:

(20 x 1 = 20)

1. LH surge precedes ovulation by
A) 12-16 hours B) 20-24 hours C) 24-36hours D) 48 hours
2. Following are features of Turners syndrome except
A) Ovarian agenesis B) Tall stature C) Cubitus valgus D) Webbing of neck
3. Ovulation can be induced by
A) Estrogen B) Progesterone C) Danazol D) Clomiphene
4. Normal pH of vagina is about
A) 1.5 B) 2.5 C) 3.5 D) 4.5
5. The commonest complication of ovarian cyst
A) Haemorrhage B) Rupture C) Torsion D) Infection
6. The best method to diagnose ulcerative cancer cervix is
A) Cervical smear B) Cervical biopsy C) Colposcopy D) None of the above
7. Corkscrew shaped glands are seen in
A) Proliferative Endometrium B) Atrophic endometrium
C) Secretory Endometrium D) Endometrial hyperplasia
8. The best diagnostic method of tubal block is
A) Ultrasonography B) Hysterosalpingography C) Colposcopy D) Hysteroscopy
9. The most common cause of hirsutism in a female is
A) Androblastoma B) Luteoma ovary
C) Adrenal hyperplasia D) PCOS
10. The following are common symptoms of endometriosis except
A) Diarrhoea B) Menorrhagia C) Dyspareunia D) Dysmenorrhoea

11. Gonococcal salpingitis is associated with all except
A) Pain B) Fever C) Vaginal bleeding D) Leukocytosis
12. The most common symptom of menopause
A) Insomnia B) Headache C) Hot flushes D) Depression
13. Nabothian follicles are seen in
A) Cervix B) Vagina C) Ovary D) Uterus
14. Commonest site of Myoma
A) Cervical B) Submucous C) Intramural D) Subserous
15. Which of the following organism causes condyloma acuminata
A) Fungus B) Virus C) Bacteria D) Protozoa
16. Granulosa and theca cell tumours are seen frequently
A) In infancy B) Before puberty C) In post-menopausal D) Reproductive age
17. The presence of pyometra in a post-menopausal woman is strongly suggestive of
A) Diabetes B) Tuberculous endometritis
C) Malignancy D) Degenerating myoma
18. The commonest ovarian neoplasm in reproductive age group is
A) Epithelial tumour B) Germ cell tumour
C) Krukenberg tumour D) Metastatic ovarian tumour
19. The treatment of acute salpingitis is
A) Antibiotics and rest B) Salpingectomy
C) Laparoscopy and fulguration D) Hysterectomy
20. Blood loss during normal menstruation is approximately
A) 15ml B) 35ml C) 75ml D) 125ml

[MBBS 0325]

THE TAMIL NADU DR. M.G.R. MEDICAL UNIVERSITY

[MBBS 0325]

MARCH 2025

Sub. Code :6086

M.B.B.S. DEGREE EXAMINATION

(For the candidates admitted from the Academic Year 2019-2020)

THIRD PROFESSIONAL MBBS PART II

PAPER II – OBSTETRICS & GYNAECOLOGY

Q.P. Code: 526086

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - C

III. Multiple Choice Questions:

(20 x 1 = 20)

1. The commonest complication of ovarian cyst
A) Haemorrhage B) Rupture C) Torsion D) Infection
2. The best method to diagnose ulcerative cancer cervix is
A) Cervical smear B) Cervical biopsy C) Colposcopy D) None of the above
3. Corkscrew shaped glands are seen in
A) Proliferative Endometrium B) Atrophic endometrium
C) Secretory Endometrium D) Endometrial hyperplasia
4. The best diagnostic method of tubal block is
A) Ultrasonography B) Hysterosalpingography C) Colposcopy D) Hysteroscopy
5. The most common cause of hirsutism in a female is
A) Androblastoma B) Luteoma ovary
C) Adrenal hyperplasia D) PCOS
6. The following are common symptoms of endometriosis except
A) Diarrhoea B) Menorrhagia C) Dyspareunia D) Dysmenorrhoea
7. Gonococcal salpingitis is associated with all except
A) Pain B) Fever C) Vaginal bleeding D) Leukocytosis
8. The most common symptom of menopause
A) Insomnia B) Headache C) Hot flushes D) Depression
9. Nabothian follicles are seen in
A) Cervix B) Vagina C) Ovary D) Uterus
10. Commonest site of Myoma
A) Cervical B) Submucous C) Intramural D) Subserous

11. Which of the following organism causes condyloma acuminata
A) Fungus B) Virus C) Bacteria D) Protozoa
12. Granulosa and theca cell tumours are seen frequently
A) In infancy B) Before puberty C) In post-menopausal D) Reproductive age
13. The presence of pyometra in a post-menopausal woman is strongly suggestive of
A) Diabetes B) Tuberculous endometritis
C) Malignancy D) Degenerating myoma
14. The commonest ovarian neoplasm in reproductive age group is
A) Epithelial tumour B) Germ cell tumour
C) Krukenberg tumour D) Metastatic ovarian tumour
15. The treatment of acute salpingitis is
A) Antibiotics and rest B) Salpingectomy
C) Laparoscopy and fulguration D) Hysterectomy
16. Blood loss during normal menstruation is approximately
A) 15ml B) 35ml C) 75ml D) 125ml
17. LH surge precedes ovulation by
A) 12-16 hours B) 20-24 hours C) 24-36hours D) 48 hours
18. Following are features of Turners syndrome except
A) Ovarian agenesis B) Tall stature C) Cubitus valgus D) Webbing of neck
19. Ovulation can be induced by
A) Estrogen B) Progesterone C) Danazol D) Clomiphene
20. Normal pH of vagina is about
A) 1.5 B) 2.5 C) 3.5 D) 4.5

[MBBS 0325]

11. The treatment of acute salpingitis is
A) Antibiotics and rest B) Salpingectomy
C) Laparoscopy and fulguration D) Hysterectomy
12. Blood loss during normal menstruation is approximately
A) 15ml B) 35ml C) 75ml D) 125ml
13. LH surge precedes ovulation by
A) 12-16 hours B) 20-24 hours C) 24-36hours D) 48 hours
14. Following are features of Turners syndrome except
A) Ovarian agenesis B) Tall stature C) Cubitus valgus D) Webbing of neck
15. Ovulation can be induced by
A) Estrogen B) Progesterone C) Danazol D) Clomiphene
16. Normal pH of vagina is about
A) 1.5 B) 2.5 C) 3.5 D) 4.5
17. The commonest complication of ovarian cyst
A) Haemorrhage B) Rupture C) Torsion D) Infection
18. The best method to diagnose ulcerative cancer cervix is
A) Cervical smear B) Cervical biopsy C) Colposcopy D) None of the above
19. Corkscrew shaped glands are seen in
A) Proliferative Endometrium B) Atrophic endometrium
C) Secretory Endometrium D) Endometrial hyperplasia
20. The best diagnostic method of tubal block is
A) Ultrasonography B) Hysterosalpingography C) Colposcopy D) Hysteroscopy

[MBBS 0325]

M.B.B.S. DEGREE EXAMINATION

(For the candidates admitted from the Academic Year 2019-2020)

THIRD PROFESSIONAL PART II SUPPLEMENTARY EXAMINATION

PAPER VI – OBSTETRICS & GYNAECOLOGY

Q.P. Code: 526086

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - A

III. Multiple Choice Questions:

(20 x 1 = 20)

1. The normal adult uterus is:
A) Retroverted and anteflexed B) Anteverted and anteflexed
C) Retroverted and retroflexed D) Anteverted and retroflexed
2. The middle layer of the uterus is called:
A) Perimetrium B) Endometrium C) Myometrium D) Mesometrium
3. The most reliable method of contraception is:
A) IUD B) OCP C) Tubal ligation D) Condom
4. OCPs primarily prevent pregnancy by:
A) Killing sperm B) Thickening cervical mucus
C) Inhibiting ovulation D) Destroying zygote
5. Which condition is not commonly associated with PCOS?
A) Obesity B) Hirsutism C) Hyperinsulinemia D) Hyperthyroidism
6. In a reproductive-age woman with sudden lower abdominal pain and positive pregnancy test, suspect:
A) Appendicitis B) Ovarian cyst C) Ectopic pregnancy D) PID
7. A molar pregnancy typically presents with:
A) Low HCG B) Bleeding and large uterus
C) Fetal heart sounds D) Normal uterine size
8. What is the gold standard diagnostic method for endometriosis?
A) Ultrasound B) CT C) Laparoscopy D) MRI
9. Treatment of choice for CIN III:
A) Repeat Pap in 6 months B) Antibiotics C) LEEP D) Observation
10. Sonohysterography is best for:
A) Cervical cancer staging B) Ovarian mass diagnosis
C) Intrauterine pathology D) Uterine perforation
11. The most common symptom of PID is:
A) Vaginal discharge B) Lower abdominal pain
C) Dyspareunia D) Menorrhagia

12. Standard treatment for PID includes:
A) Antifungals B) Broad-spectrum antibiotics C) NSAIDs D) Antivirals
13. Most appropriate investigation in a 45-year-old woman with AUB:
A) Pap smear B) Pregnancy test C) Endometrial biopsy D) FSH level
14. First-line management of AUB due to anovulation in adolescents:
A) Hysterectomy B) Iron supplements
C) Combined oral contraceptives D) D&C
15. Which gynecologic malignancy has the worst prognosis?
A) Cervical B) Endometrial C) Ovarian D) Vulvar
16. The most common risk factor for endometrial cancer is:
A) Early menopause B) Obesity
C) Smoking D) Multiple pregnancies
17. The most common location of fibroid is:
A) Submucosal B) Subserosal C) Intramural D) Pedunculated
18. Letrozole is a/an:
A) GnRH agonist B) Aromatase inhibitor
C) Estrogen agonist D) Selective estrogen receptor modulator
19. Ulipristal acetate as emergency contraception is effective up to:
A) 24 hours B) 48 hours C) 72 hours D) 120 hours
20. Intraoperative ureteral injury is most commonly associated with:
A) Myomectomy B) Tubal ligation
C) Hysterectomy D) Ovarian cystectomy

[MBBS 0625]

12. Letrozole is a/an:
A) GnRH agonist B) Aromatase inhibitor
C) Estrogen agonist D) Selective estrogen receptor modulator
13. Ulipristal acetate as emergency contraception is effective up to:
A) 24 hours B) 48 hours C) 72 hours D) 120 hours
14. Intraoperative ureteral injury is most commonly associated with:
A) Myomectomy B) Tubal ligation
C) Hysterectomy D) Ovarian cystectomy
15. The normal adult uterus is:
A) Retroverted and anteflexed B) Anteverted and anteflexed
C) Retroverted and retroflexed D) Anteverted and retroflexed
16. The middle layer of the uterus is called:
A) Perimetrium B) Endometrium C) Myometrium D) Mesometrium
17. The most reliable method of contraception is:
A) IUD B) OCP C) Tubal ligation D) Condom
18. OCPs primarily prevent pregnancy by:
A) Killing sperm B) Thickening cervical mucus
C) Inhibiting ovulation D) Destroying zygote
19. Which condition is not commonly associated with PCOS?
A) Obesity B) Hirsutism C) Hyperinsulinemia D) Hyperthyroidism
20. In a reproductive-age woman with sudden lower abdominal pain and positive pregnancy test, suspect:
A) Appendicitis B) Ovarian cyst C) Ectopic pregnancy D) PID

[MBBS 0625]

M.B.B.S. DEGREE EXAMINATION

(For the candidates admitted from the Academic Year 2019-2020)

THIRD PROFESSIONAL PART II SUPPLEMENTARY EXAMINATION

PAPER VI – OBSTETRICS & GYNAECOLOGY

Q.P. Code: 526086

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - C

III. Multiple Choice Questions:

(20 x 1 = 20)

1. First-line management of AUB due to anovulation in adolescents:
A) Hysterectomy B) Iron supplements
C) Combined oral contraceptives D) D&C
2. Which gynecologic malignancy has the worst prognosis?
A) Cervical B) Endometrial C) Ovarian D) Vulvar
3. The most common risk factor for endometrial cancer is:
A) Early menopause B) Obesity
C) Smoking D) Multiple pregnancies
4. The most common location of fibroid is:
A) Submucosal B) Subserosal C) Intramural D) Pedunculated
5. Letrozole is a/an:
A) GnRH agonist B) Aromatase inhibitor
C) Estrogen agonist D) Selective estrogen receptor modulator
6. Ulipristal acetate as emergency contraception is effective up to:
A) 24 hours B) 48 hours C) 72 hours D) 120 hours
7. Intraoperative ureteral injury is most commonly associated with:
A) Myomectomy B) Tubal ligation
C) Hysterectomy D) Ovarian cystectomy
8. The normal adult uterus is:
A) Retroverted and anteflexed B) Anteverted and anteflexed
C) Retroverted and retroflexed D) Anteverted and retroflexed
9. The middle layer of the uterus is called:
A) Perimetrium B) Endometrium C) Myometrium D) Mesometrium
10. The most reliable method of contraception is:
A) IUD B) OCP C) Tubal ligation D) Condom
11. OCPs primarily prevent pregnancy by:
A) Killing sperm B) Thickening cervical mucus
C) Inhibiting ovulation D) Destroying zygote

12. Which condition is not commonly associated with PCOS?
A) Obesity B) Hirsutism C) Hyperinsulinemia D) Hyperthyroidism
13. In a reproductive-age woman with sudden lower abdominal pain and positive pregnancy test, suspect:
A) Appendicitis B) Ovarian cyst C) Ectopic pregnancy D) PID
14. A molar pregnancy typically presents with:
A) Low HCG B) Bleeding and large uterus
C) Fetal heart sounds D) Normal uterine size
15. What is the gold standard diagnostic method for endometriosis?
A) Ultrasound B) CT C) Laparoscopy D) MRI
16. Treatment of choice for CIN III:
A) Repeat Pap in 6 months B) Antibiotics C) LEEP D) Observation
17. Sonohysterography is best for:
A) Cervical cancer staging B) Ovarian mass diagnosis
C) Intrauterine pathology D) Uterine perforation
18. The most common symptom of PID is:
A) Vaginal discharge B) Lower abdominal pain
C) Dyspareunia D) Menorrhagia
19. Standard treatment for PID includes:
A) Antifungals B) Broad-spectrum antibiotics C) NSAIDs D) Antivirals
20. Most appropriate investigation in a 45-year-old woman with AUB:
A) Pap smear B) Pregnancy test C) Endometrial biopsy D) FSH level

[MBBS 0625]

M.B.B.S. DEGREE EXAMINATION

(For the candidates admitted from the Academic Year 2019-2020)

THIRD PROFESSIONAL PART II SUPPLEMENTARY EXAMINATION

PAPER VI – OBSTETRICS & GYNAECOLOGY

Q.P. Code: 526086

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - D

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Letrozole is a/an:
A) GnRH agonist B) Aromatase inhibitor
C) Estrogen agonist D) Selective estrogen receptor modulator
2. Ulipristal acetate as emergency contraception is effective up to:
A) 24 hours B) 48 hours C) 72 hours D) 120 hours
3. Intraoperative ureteral injury is most commonly associated with:
A) Myomectomy B) Tubal ligation
C) Hysterectomy D) Ovarian cystectomy
4. The normal adult uterus is:
A) Retroverted and anteflexed B) Anteverted and anteflexed
C) Retroverted and retroflexed D) Anteverted and retroflexed
5. The middle layer of the uterus is called:
A) Perimetrium B) Endometrium C) Myometrium D) Mesometrium
6. The most reliable method of contraception is:
A) IUD B) OCP C) Tubal ligation D) Condom
7. OCPs primarily prevent pregnancy by:
A) Killing sperm B) Thickening cervical mucus
C) Inhibiting ovulation D) Destroying zygote
8. Which condition is not commonly associated with PCOS?
A) Obesity B) Hirsutism C) Hyperinsulinemia D) Hyperthyroidism
9. In a reproductive-age woman with sudden lower abdominal pain and positive pregnancy test, suspect:
A) Appendicitis B) Ovarian cyst C) Ectopic pregnancy D) PID
10. A molar pregnancy typically presents with:
A) Low HCG B) Bleeding and large uterus
C) Fetal heart sounds D) Normal uterine size
11. What is the gold standard diagnostic method for endometriosis?
A) Ultrasound B) CT C) Laparoscopy D) MRI

12. Treatment of choice for CIN III:
A) Repeat Pap in 6 months B) Antibiotics C) LEEP D) Observation
13. Sonohysterography is best for:
A) Cervical cancer staging B) Ovarian mass diagnosis
C) Intrauterine pathology D) Uterine perforation
14. The most common symptom of PID is:
A) Vaginal discharge B) Lower abdominal pain
C) Dyspareunia D) Menorrhagia
15. Standard treatment for PID includes:
A) Antifungals B) Broad-spectrum antibiotics C) NSAIDs D) Antivirals
16. Most appropriate investigation in a 45-year-old woman with AUB:
A) Pap smear B) Pregnancy test C) Endometrial biopsy D) FSH level
17. First-line management of AUB due to anovulation in adolescents:
A) Hysterectomy B) Iron supplements
C) Combined oral contraceptives D) D&C
18. Which gynecologic malignancy has the worst prognosis?
A) Cervical B) Endometrial C) Ovarian D) Vulvar
19. The most common risk factor for endometrial cancer is:
A) Early menopause B) Obesity
C) Smoking D) Multiple pregnancies
20. The most common location of fibroid is:
A) Submucosal B) Subserosal C) Intramural D) Pedunculated

[MBBS 0625]

10. The following are the primary sites of acute gonococcal infection except:
A) Urethra
B) Bartholin's gland
C) Skene's gland (paraurethral glands)
D) Ectocervix
11. All of the following support uterus except:
A) Cardinal ligament
B) Round ligament
C) Pubocervical ligament
D) Mackenrodt's ligament
12. Trauma during child birth is a risk factor for:
A) Endometriosis
B) Prolapse
C) Abortion
D) PID
13. Liquefaction time of semen is:
A) 30 mins
B) 60 mins
C) 50 mins
D) 10 minutes
14. Ring of fire appearance in USG
A) Chocolate cyst of ovary
B) Ectopic pregnancy
C) Hydatidiform mole
D) Poly cystic ovary
15. All of the following are the indications for myomectomy in a case of fibroid uterus except:
A) Associated infertility
B) Recurrent pregnancy loss
C) Pressure symptoms
D) Red degeneration
16. Most common accepted theory for endometriosis is:
A) Implantation theory
B) Metastatic theory
C) Coelomic metaplasia
D) Histogenesis theory
17. Causative organism of chancroid
A) Chlamydia trachomatis
B) Treponema pallidum
C) Hemophilus ducreyi
D) Neisseria gonorrhoea
18. Endometrial hyperplasia is often associated with:
A) Cystic teratoma
B) Endodermal tumor
C) PCOS
D) Sertoli cell tumour
19. Posterior colpotomy is done for:
A) Pelvic hematocoele
B) Pelvic abscess
C) Ovarian abscess
D) All
20. Cause of secondary dysmenorrhea in a young female:
A) Tuberculosis
B) CIN
C) Endometriosis
D) Subserosal fibroid

M.B.B.S. DEGREE EXAMINATION
(For the candidates admitted from the Academic Year
2019-2020, 2020-2021 & 2021-2022)

THIRD PROFESSIONAL PART II
PAPER II – OBSTETRICS & GYNAECOLOGY

Q.P. Code: 526086

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - B

III. Multiple Choice Questions:

(20 x 1 = 20)

1. All of the following are the indications for myomectomy in a case of fibroid uterus except:
A) Associated infertility
B) Recurrent pregnancy loss
C) Pressure symptoms
D) Red degeneration
2. Most common accepted theory for endometriosis is:
A) Implantation theory
B) Metastatic theory
C) Coelomic metaplasia
D) Histogenesis theory
3. Causative organism of chancroid
A) Chlamydia trachomatis
B) Treponema pallidum
C) Hemophilus ducreyi
D) Neisseria gonorrhoea
4. Endometrial hyperplasia is often associated with:
A) Cystic teratoma
B) Endodermal tumor
C) PCOS
D) Sertoli cell tumour
5. Posterior colpotomy is done for:
A) Pelvic hematocele
B) Pelvic abscess
C) Ovarian abscess
D) All
6. Cause of secondary dysmenorrhea in a young female:
A) Tuberculosis
B) CIN
C) Endometriosis
D) Subserosal fibroid
7. For hormonal study sample should be taken from which wall of vagina
A) Anterior wall
B) Posterior wall
C) Lateral wall
D) Any wall
8. Just before ovulation there is:
A) LH surge
B) FSH surge
C) LH and FSH surge
D) None

9. Luteal phase defect is due to
A) Uterine scarring
B) Chromosomal abnormality
C) Abnormal release of GNRH
D) Absence of Oocytes in ovaries
10. Menopause transition of a female is from
A) 30-49 years
B) 35-49 years
C) 40-51 years
D) 40-60 years
11. Which Hormone level will be raised in menopause?
A) Progesterone
B) FSH
C) ACTH
D) Estradiol
12. Drug used to treat post-menopausal osteopenia Except
A) Alendronate
B) Strontium
C) Calcitonin
D) Hydroxychloroquine
13. Most common cause of Hirsutism in a young female:
A) Ovarian tumor
B) Adrenal tumor
C) CAH
D) PCOS
14. Unicollis bicorn is means:
A) Two uterine cavity with one cervix
B) Single vagina with double uterus
C) Complete septum of uterus
D) Double uterus and double cervix
15. Drugs used for emergency contraception
A) Norethisterone
B) Dydrogesterone
C) Levonorgestrel
D) Medroxyprogesterone acetate
16. The following are the primary sites of acute gonococcal infection except:
A) Urethra
B) Bartholin's gland
C) Skene's gland (paraurethral glands)
D) Ectocervix
17. All of the following support uterus except:
A) Cardinal ligament
B) Round ligament
C) Pubocervical ligament
D) Mackenrodt's ligament
18. Trauma during child birth is a risk factor for:
A) Endometriosis
B) Prolapse
C) Abortion
D) PID
19. Liquefaction time of semen is:
A) 30 mins
B) 60 mins
C) 50 mins
D) 10 minutes
20. Ring of fire appearance in USG
A) Chocolate cyst of ovary
B) Ectopic pregnancy
C) Hydatidiform mole
D) Poly cystic ovary

M.B.B.S. DEGREE EXAMINATION
(For the candidates admitted from the Academic Year
2019-2020, 2020-2021 & 2021-2022)

THIRD PROFESSIONAL PART II
PAPER II – OBSTETRICS & GYNAECOLOGY

Q.P. Code: 526086

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - C

III. Multiple Choice Questions:

(20 x 1 = 20)

1. The following are the primary sites of acute gonococcal infection except:
A) Urethra
B) Bartholin's gland
C) Skene's gland (paraurethral glands)
D) Ectocervix
2. All of the following support uterus except:
A) Cardinal ligament
B) Round ligament
C) Pubocervical ligament
D) Mackenrodt's ligament
3. Trauma during child birth is a risk factor for:
A) Endometriosis
B) Prolapse
C) Abortion
D) PID
4. Liquefaction time of semen is:
A) 30 mins
B) 60 mins
C) 50 mins
D) 10 minutes
5. Ring of fire appearance in USG
A) Chocolate cyst of ovary
B) Ectopic pregnancy
C) Hydatidiform mole
D) Poly cystic ovary
6. All of the following are the indications for myomectomy in a case of fibroid uterus except:
A) Associated infertility
B) Recurrent pregnancy loss
C) Pressure symptoms
D) Red degeneration
7. Most common accepted theory for endometriosis is:
A) Implantation theory
B) Metastatic theory
C) Coelomic metaplasia
D) Histogenesis theory
8. Causative organism of chancroid
A) Chlamydia trachomatis
B) Treponema pallidum
C) Hemophilus ducreyi
D) Neisseria gonorrhoea

9. Endometrial hyperplasia is often associated with:
A) Cystic teratoma
B) Endodermal tumor
C) PCOS
D) Sertoli cell tumour
10. Posterior colpotomy is done for:
A) Pelvic hematocele
B) Pelvic abscess
C) Ovarian abscess
D) All
11. Cause of secondary dysmenorrhea in a young female:
A) Tuberculosis
B) CIN
C) Endometriosis
D) Subserosal fibroid
12. For hormonal study sample should be taken from which wall of vagina
A) Anterior wall
B) Posterior wall
C) Lateral wall
D) Any wall
13. Just before ovulation there is:
A) LH surge
B) FSH surge
C) LH and FSH surge
D) None
14. Luteal phase defect is due to
A) Uterine scarring
B) Chromosomal abnormality
C) Abnormal release of GnRH
D) Absence of Oocytes in ovaries
15. Menopause transition of a female is from
A) 30-49 years
B) 35-49 years
C) 40-51 years
D) 40-60 years
16. Which Hormone level will be raised in menopause?
A) Progesterone
B) FSH
C) ACTH
D) Estradiol
17. Drug used to treat post-menopausal osteopenia Except
A) Alendronate
B) Strontium
C) Calcitonin
D) Hydroxychloroquine
18. Most common cause of Hirsutism in a young female:
A) Ovarian tumor
B) Adrenal tumor
C) CAH
D) PCOS
19. Unicorn uterus is means:
A) Two uterine cavity with one cervix
B) Single vagina with double uterus
C) Complete septum of uterus
D) Double uterus and double cervix
20. Drugs used for emergency contraception
A) Norethisterone
B) Dydrogesterone
C) Levonorgestrel
D) Medroxyprogesterone acetate

THE TAMIL NADU Dr. M.G.R. MEDICAL UNIVERSITY

[MBBS 0326]

MARCH 2026

Sub. Code :6086

M.B.B.S. DEGREE EXAMINATION
(For the candidates admitted from the Academic Year
2019-2020, 2020-2021 & 2021-2022)

THIRD PROFESSIONAL PART II
PAPER II – OBSTETRICS & GYNAECOLOGY

Q.P. Code: 526086

Time: 20 Minutes

Maximum : 20 Marks

Answer All Questions

Shade the correct answer with Black or Blue ink ball point pen

SET - D

III. Multiple Choice Questions:

(20 x 1 = 20)

1. Which Hormone level will be raised in menopause?
A) Progesterone B) FSH C) ACTH D) Estradiol
2. Drug used to treat post-menopausal osteopenia Except
A) Alendronate B) Strontium
C) Calcitonin D) Hydroxychloroquine
3. Most common cause of Hirsutism in a young female:
A) Ovarian tumor B) Adrenal tumor
C) CAH D) PCOS
4. Unicollis bicorn is means:
A) Two uterine cavity with one cervix B) Single vagina with double uterus
C) Complete septum of uterus D) Double uterus and double cervix
5. Drugs used for emergency contraception
A) Norethisterone B) Dydrogesterone
C) Levonorgestrel D) Medroxyprogesterone acetate
6. The following are the primary sites of acute gonococcal infection except:
A) Urethra B) Bartholin's gland
C) Skene's gland (paraurethral glands) D) Ectocervix
7. All of the following support uterus except:
A) Cardinal ligament B) Round ligament
C) Pubocervical ligament D) Mackenrodt's ligament
8. Trauma during child birth is a risk factor for:
A) Endometriosis B) Prolapse
C) Abortion D) PID
9. Liquefaction time of semen is:
A) 30 mins B) 60 mins C) 50 mins D) 10 minutes

10. Ring of fire appearance in USG
A) Chocolate cyst of ovary
B) Ectopic pregnancy
C) Hydatidiform mole
D) Poly cystic ovary
11. All of the following are the indications for myomectomy in a case of fibroid uterus except:
A) Associated infertility
B) Recurrent pregnancy loss
C) Pressure symptoms
D) Red degeneration
12. Most common accepted theory for endometriosis is:
A) Implantation theory
B) Metastatic theory
C) Coelomic metaplasia
D) Histogenesis theory
13. Causative organism of chancroid
A) Chlamydia trachomatis
B) Treponema pallidum
C) Hemophilus ducreyi
D) Neisseria gonorrhoea
14. Endometrial hyperplasia is often associated with:
A) Cystic teratoma
B) Endodermal tumor
C) PCOS
D) Sertoli cell tumour
15. Posterior colpotomy is done for:
A) Pelvic hematocele
B) Pelvic abscess
C) Ovarian abscess
D) All
16. Cause of secondary dysmenorrhea in a young female:
A) Tuberculosis
B) CIN
C) Endometriosis
D) Subserosal fibroid
17. For hormonal study sample should be taken from which wall of vagina
A) Anterior wall
B) Posterior wall
C) Lateral wall
D) Any wall
18. Just before ovulation there is:
A) LH surge
B) FSH surge
C) LH and FSH surge
D) None
19. Luteal phase defect is due to
A) Uterine scarring
B) Chromosomal abnormality
C) Abnormal release of GnRH
D) Absence of Oocytes in ovaries
20. Menopause transition of a female is from
A) 30-49 years
B) 35-49 years
C) 40-51 years
D) 40-60 years
